

USDA Foreign Agricultural Service

GAIN Report

Global Agricultural Information Network

THIS REPORT CONTAINS ASSESSMENTS OF COMMODITY AND TRADE ISSUES MADE BY USDA STAFF AND NOT NECESSARILY STATEMENTS OF OFFICIAL U.S. GOVERNMENT POLICY

Voluntary Public

Date: 4/25/2012

GAIN Report Number: RS1227

Russian Federation

Post: Moscow

CU Draft on New MRLs for Pesticides in Agricultural Products

Report Categories:

Sanitary/Phytosanitary/Food Safety

Approved By:

Levin Flake

Prepared By:

Yelena Vassilieva, Marina Muran

Report Highlights:

On April 10, 2012 the Customs Union of the Republic of Belarus, Republic of Kazakhstan and Russian Federation (CU) issued for public discussion the draft on Amendments to the Unified Sanitary-Epidemiological and Hygiene Requirements for Commodities Subject to Sanitary-Epidemiological Surveillance (Control). Part of the proposed amendments concern Maximum Residue Limits (MRLs) for pesticides and chemicals in external entities, including agricultural products. The proposed new edition of the MRLs for agricultural products is given in the report. This draft is open for public comment for 60 days from April 10th, 2012.

General Information:

On April 10, 2012, the Euro-Asian Economic Commission, which replaced the Customs Union Commission of the Republic of Belarus, Republic of Kazakhstan and Russian Federation ^[i], issued for public discussion the draft on Amendments to the Unified Sanitary-Epidemiological and Hygiene Requirements for Commodities Subject to Sanitary-Epidemiological Surveillance (Control): <http://tsouz.ru/db/techregulation/Documents/Решение%20САН%20требования.pdf> . The amendments concern several issues, including new Maximum Residue Limits (MRLs) for pesticides and chemicals in external entities, including agricultural products. The current Customs Union MRLs for pesticides include a list of almost 490 chemicals (active substances) with specified maximum allowed levels in the human body, in soil, in reservoir water, in working air, in open air, and in products. FAS/Moscow reported on the MRLs for pesticides in Agricultural Products adopted by the CU decision 342 of August 18, 2010 in the GAIN Report RS1076 Customs Union Update on MRLs for Pesticides in Ag Products _ Moscow_Russian_Federation_12-22-2010.doc. The draft of new MRLs for pesticides in agricultural products is in the Annex below. (**NOTE:** in the Annex below the proposed amendments and/or additions, compared with what has been in force since August 2010, are marked in **YELLOW**; text **IN RED** with strike through means that this text is deleted or replaced in the proposed new draft.)

Compared with the current regulations, the draft differs in the following areas:

- The draft includes 77 more chemical ingredients, taking the total number of chemical ingredients in the draft to 563 compared with 486 in the current legislation;
- For many agricultural plant products MRLs of pesticides are higher (more liberal) than in the current requirements;
- Agricultural products are divided by many subgroups, and for many chemical ingredients the requirements are determined for each small sub-group. For example, if in the current regulations the MRL for a certain ingredient is set for the whole group of berries, in the proposed draft the MRLs are set for each separate berry;
- The draft includes MRLs for many pesticides in animal products, such as meat of cattle, goat, pig, sheep, poultry, milk and offal of these animals. In the current legislation only a few MRLs are given for animal products, and the animal products are not divided by specific types.

There is a 60-day public comment period, starting April 10, 2012. Interested U.S. parties are encouraged to share their comments and concerns with USDA.

^[i] For more information on the status of the Eurasian Economic Commission (EEC), and its functions as a successor of the Custom Union Commission see GAIN RS1212 Customs Union Ag Times No. 1_Moscow_Russian Federation_21.02.2012.doc

Annex:

NOTE: The Table below is an extract from the draft **Hygiene Norms of for Chemicals and Pesticides in External Entities, in Agricultural Raw Material and in Food Products** and shows only names of chemicals and MRLs in agricultural and food products. Please note that this table is an unofficial translation, and for more detail refer to the document itself:

<http://tsouz.ru/db/techregulation/Documents/Решение%20САН%20требования.pdf>.

Abbreviations and symbols used in the table:

- MPL – maximum permissible level;
- TMPL - Temporary maximum permissible level marked with asterisk (*);
- MPL for imported production is marked with two asterisks (**);
- TMPL for imported products is marked with two asterisks (**);
- NR - substance not rated in the given media;
- RNR - substance not required in the given media;
- CATTLE - cattle

Table 1. MRL/TMPL in Agricultural Products (mg/kg)

NO	Name of active ingredient (Column 2)	MPL/TMPL in product (mg/kg) (Column 8)
1	β -digidroheptachlore	Potatoes, cottonseed (oil), grapes- 0.15; sugar beet, vegetables (except potatoes) - 0.2; blue poppy -0.15*
2	(indolyl -3) acetic acid	RNR
3	(chloride-N, N- dimethyl -N-)-(2-chloroethyl) hydrozinia	Cereal grain, fruits (pomaceous fruits), potatoes - NR
4	0-(2, 4- dichloro phenol)-S- propyl - O-ethylphosfate	Fruits (pomaceous fruits, stone fruits), citrus fruits (pulp), cabbage, potatoes, meat - 0.01; grapes, berries -0.01*; cottonseed (oil) – 0.02*; sunflower (seeds) – 0. 1 *; sugar beet - 0.02
5	0-(4- tert-butyl -2-chlorophenyl) -0-methyl -N- methyl- amidophosphate	Meat, meat products - 0.3
6	0-methyl-0-(2, 4, 5- trichlorophenil) -0-ethyltiophosphate	Cucumbers, tomatoes, sugar beet, cabbage, fruits (pomaceous fruits, stone fruits), grapes, mushrooms -1.0; tobacco - 0.7; citrus fruits (pulp)- 0.3*; tea - 0.5; cottonseed (oil) -0. 1
7	0-ethyl-0- phenyl-S- propilthiophosphate	All food products –NR
8	0,0-Dimethyl-0-(4-methylthio-3-methyl-phenil) thiophosphate	NR
9	1,1-di-(4-chlore- phenil) - 2,2,2-trichloroethane (DDT)	Grain of cereals – 0.1*, meat of mammals, except sea animals – 5.0, poultry meat (fresh, cooled and frozen) -0.3; eggs – 0.1; milk – 0.02*, carrots – 0.2*, **, byproducts (liver,

		<p>kidneys), sausages, cookery, canned meat and poultry – as per raw material (in terms of fat); eggs, flax (seeds), rape (seeds), mustard, vegetables, melons, mushrooms, potatoes, fruits, berries, grapes, vegetable oil, deodorized, of best purification, gelatin - 0,1; milk and cultured milk products, legumes, soya (beans) - 0,05; milk processing products (cheeses, curd products, butter, cream, sour cream), concentrates of milk, whey proteins, dry milk and milk products (in terms of fat), animal fat - 1.0; freshwater fish (fresh, cooled, frozen)-0.3; sea tuna fish, (fresh, cooled, frozen), meat of sea animals, non-deodorized vegetable oil, fish fat - 0.2; fish: salty, smoked, sun-cared - 0.4; fish cans (freshwater, seawater, tuna fishes, meat of sea animals) – as to raw material; liver of fishes and products made of it - 3.0; caviar, sturgeons, salmons, fat herring - 2.0; cereal grain, corn - 0.02; flour confectionery – 0.02; starch and syrup made of corn-0.05; starch and syrup made of potatoes-0.1; flour, cereals - as per; seeds of sunflower, peanut, nuts, cocoa (beans), cocoa-products - 0,15; fruit and vegetables cans- as per raw material; juices - as per raw material; honey - 0.005; tobacco -0.7; protein products made of seeds of cereals, leguminous plants etc., - 0.01; Baby products: adapted milk mixes (for children 0—3 months)-0.01; products for children 4-12 months: milk – 0.01, cottage cheese 18% - 0.06, meat, cereals -0.01; vegetables, potatoes, fruits - 0.005; butter - 0.2; vegetable oil - 0.1</p>
10	1,1-dioxotiolanin-3- three ethylene salt of dithiocarbarnic acid	NR
11	1- (2-chloreetoxicarbonilmethyl)-calcium naphthalene sulfoacids	Potatoes -NR
12	[1-(4-nitrophenyl) -2- amino -1,3-propandiol] nitrate	Tomatoes — NR
13	2, 3, 6-TBA	Wheat -0.05*
14	2, 4-D acid	cereal grain – 2.0, millet, corn (grain) - 0.05;
15	2, 4-D butyl ether	sorghum – 0.01*, **; corn (oil)-0.1; milk-0.01*
16	2, 4-D low-volatile esters +2,4D 2-ethylhexyl ester	0.04; butter-0.1; flour, grits – as per raw material*; fresh water fish -0.01*; citrus fruits
17	2, 4-D octyl ester	-1.0**; berries and other small fruits, milled

		rice – 0.1 *, **; mammals' sub-products – 5.0*, **; eggs, seed type fruits, soya (beans) – 0.01* **; meat - 0.08 of mammals, except sea animals, potato tree nuts – 0.2*, **; poultry meat and sub-products, stone type fruits, sugar cane, corn sweet, table (boiled in cobs) – 0.05*, **
18	2, 4-DB	Cereal grain – NR
19	2- amino -6- dimethylamino -4-chloride-1 ,3,5- triazine (metabolite and preproduct of gramex synthesis)	NR
20	2-carbometoxi-amino-qunazon	NR
21	2-methyl-4-dimethylaminomethyl-benzimidazole -5-ole dihydropochloride	Corn, cucumbers –NR
22	2-methyl-4-oxo-3-(prop-2-enyl)-2-cyclopenten-2- en-1-il-2,2-dimethyl-3-(2-methyl- prop -1-enyl- cyclopropancarbonat	NR
23	2-oxo-2,5-dihydrofueran	Cereal grain, corn (grain), rice -0,2; cucumbers, eabbage NR
24	2-phenylphenol	Citrus fruits – 10.0 *, **; dried soft part of citrus fruits – 60.0*, **; orange juice – 0.5*, **; seed type fruits – 20.0 *, **
25	2-chloreathylphosphon acid benzimidazol sault	Potatoes NR
26	2-(diphenylacetyl)1H-inden-1,3-2H- dion	NR
27	2-[4-(1-methylethyl) phenyl phenylacetyl]-1H-indan-1,3 dion	NR
28	2-[(4-chlorophenil) phenilacetyl]-1H-inden-1,3 (2H) -dion	NR
29	3,3-dichlore-tri-cyclo-(2,2,1)-hepta-5-en-2-spiro-[2'-(4',5-dichlore-4'-cyclopenten-1',3'-dion]	NR
30	5-ethyl-5-hydroximethyl-2-(phuril-2)-1,3- dioxane	Cereal grain - 0,1; pepper, tomatoes-0.05
31	5,6,7- trichloro -3- benzothiadiazine -oxide -1	Sugar beet -0.04
32	6-methyl-2- thiouracil sodium salt	Panicum, oats – NR
33	Bacillus thuringiensis , var. dendrolimus (sporo – crystalline complex and ectotoxin	RNR
34	Bacillus thuringiensis, var. insektus (sporo – crystalline complex and ectotoxin)	RNR
35	Bacillus thuringiensis, var. kurstaki (sporo – crystalline complex)	RNR

36	Bacillus thuringiensis, var. tenebrionis (sporo -- crystalline complex and ectotoxin	RNR
37	Bacillus thuringiensis, var. thuringiensis (sporo - crystalline complex)	RNR
38	Bacillus thuringiensis, var. thuringiensis (sporo -- crystalline complex and ectotoxin	RNR
39	Beaveria bassiana (conidia)	RNR
40	EPTC	Corn (grain), vegetable oil, sugar beet-0.05
41	MCPA	Pea, millet, rice, potatoes, sunflower (oil), cereal grain-0.05, beans-0.1
42	MCPB	Cereal grain, legumes-0.1
43	N-hexyloxymethylazepin	NR
44	NN-β- oxyethyl (morpholiny chloride)	Cottonseed (oil), buckwheat - NR
45	N,N - dimethyl - N '(3-chlorephenil) guanidine	Cucumbers- 1.0
46	N- β - methoxy -ethylchloraceto-0-toluide	Cottonseed (oil) -0.25; corn -0.5*
47	N- β -etoxiethylchloreacetamid	NR
48	N-(isopropoxi -carbonil-0-(4-chlorophenilcarbamoila)- ethanolamine	All food products - NR
49	N-(4-chlorophenyl) -4, 6-dimethyl-3-carboxipiri-din-2-on	Cottonseed (oil) - NR
50	N-methyl-0-tolilcarbamat	Milk, milk products, eggs - NR
51	-2, 6-lutidine M- oxide	Tomatoes, cucumbers -0.04;
52	S-methyl-N-methyl- carbomoil) oxitiaceti-midat	NR
53	Pseudomonas syringae (bacteriophage)	RNR
54	Verticillium lecanii (conidin)	RNR
55	Abamectin	Hops (dry) - 0.1*,**' nuts (almonds, walnut) - 0.01*,**; almonds (in shell) - 0.1*,**; pomaceous fruits, tomatoes - 0.02; citrus fruits - 0.01*,**; cucumbers-0.01; leaf lettuce - 0.05*,**; cottonseed (seeds) - 0.01*,**; melons, water melons, pumpkins - 0.01*,**; pepper Chile (dry) - 0.2*,**, strawberry, sweet pepper (including pod pepper) - 0.02*,**, sub-products (goat), fat, liver (C, goats) - 0.01*,**; eggplants - 0.01; grapes - 0.01 tomatoes; pepper, eggplants, grapes - 0.003
56	Aversectin C	Cucumbers, tomatoes, potatoes, fruits (pomaceous fruits), currant -0.005; meat-0.004; offal-0.01; fat-0.024; milk-0.001
57	Azimsulfuron	Rice-0.02
58	Azinphos-methyl	Nuts: pecan, walnuts - 0.3*,**, almonds - 0.05*,**; almonds in shell - 5,0*,**; pomaceous fruits: apples - 0.05*,**, pears -

		2.0*,**; stone type fruits (except prunes)- 2.0*,**; berries: blueberry – 5.0*,**, cranberry – 0.1*,**; broccoli, fruits (except listed above), sweet pepper, tomatoes – 1.0*,**; cottonseed (seeds), melons, water-melons, sugar cane – 2.0*,**; pepper Chili (dry) – 10.0*,**; potatoes, soya (dry beans) – 0.05*,**; vegetables (except listed above) – 0.5*,**
59	Aziprotrin	Vegetables (except potatoes) - 0.2
60	Azoxistrobin	Artichoke, cabbage (all types), celery, rice, berries and other small fruits (except cranberry, grapes and strawberry) – 5.0*,**; asparagus, tree nuts (except pistachios) – 0.01*,**; pistachios – 1.0*,**; almonds in shell – 7.0*,**; bananas – stone type fruits – 2.0*,** - grapes – 0.2, cereal grain: barley, oats – 0.5, wheat, rye, triticale – 0.3; soya (beans), sunflowerseed (seeds), cranberry – 0.5*,**; vegetables with bulbs fit for human consumption (except onion), strawberry – 10.0*,**; onion – 10.0; citrus fruits – 15.0*,**; cottonseed (seeds), mango – 0.7*,**; fruit-bearing vegetables (except tomatoes, pumpkins, cucumbers), legumes, lettuce (leaf and loaf) – 3.0*,**; tomatoes, cucumbers – 0.3*,** -2.0; pumpkin, vegetables with edible roots – 1.0*,**; potatoes – 0.05; hops (dry), pepper Chili (dry) – 30.0*,**; corn (grain) – 0.02*,**; corn (oil) – 1.0*,**, papaya, chicory - 0.3*,**; peanuts – 0.2*,**; milk, eggs, poultry meat, offal (poultry) – 0.01*,**; meat of mammal animals (except sea animals) – 0.05*,**; milk fat – 0.03*,**; offal of mammal animals – 0.7*,** onion – 0.05,-, cucumbers – 0.2
61	Azocyclotin	Pomaceous fruits – 0.2*,**, currant (red, white, black) – 0.1*,**, grapes – 0.3*,**; oranges (including hybrids) – 0.2*,**
62	Acvo-N-oxi-2-methylpiridin manganese (II) chloride	Cereal grain - 0.08
63	Akrinatriin	Fruits (pomaceous fruits) - 0.03*
64	Acraldehyde	RNR
65	Alachlor	Soya (beans, oil), corn (grain) -0.02*
66	Aldrin and dueldrin	Vegetables with edible bulbs (garlic, onion, etc.), citrus fruits, leaf vegetables (lettuce, spinach, parsley etc.), legumes (beans, peas),

		pomaceous fruits, pulses – 0.05*,**, cereal grain – 0.02*,**, pumpkin type, vegetables with edible roots – 0.1*,**, potato, beets – 0.1 – 0.002 ; peas (juicy seeds) – 1.0*,**; meat of mammals (except sea mammals), poultry meat – 0.2*,**, milk – 0.006*; eggs – 0.1*,**, cabbage-0.004; vine, products of vegetables processing-0.005; animal fat, milk, cream, curd-0.04; sugar-0.02
67	Aldicarb	Soya (beans), cereal grain – 0.02*,**, beans, Brussels cabbage, coffee beans, cottonseed (seeds), onion, sorghum, sugar cane, sweet potato – 0.1*,**, citrus fruits, grapes – 0.2*,**, corn, sugar beet, sunflowerseed (seeds) – 0.05*,**, peanuts – 0.02*,**, vegetable oil for food consumption (cottonseed, peanut) – 0.01*,**, pecan nut – 1.0*,**, meat of mammals (except sea mammals) – 0.01*,**, milk – 0.01*,**
68	Alkyl - ether - sodium salt sulfate	NR
69	Alloxidim natrium	Sugar beet, red beet - 0.05
70	Alfa Cypermethrin (mixture of Cypermethrin isomers)	Grapes, fruits (stone fruits), red beets, mustard, tomatoes, wild-growing mushrooms and berries – 0.005; pea-0.1; rapeseed (seeds, oil), cereal grains, potatoes, sugar beets, fruits (pomaceous fruits); corn (grains, oil)-0.05
71	Aluminum fosethyl	Grapes - 0.8; cucumbers - 0.5; onion -0.01; dry hop - 1.0; tomatoes – 100.0; cucumbers – 75.0
72	Amidosulphuron	Cereal grain -0.1 ; corn (grain, oil) -0.5
73	Free amino acids	RNR
74	Aminopirialid	Cereal grain (barley, wheat, oat), triticale- 0.1; offal of mammals – 0.05*,**, eggs – 0.01*,**, kidney of C, goats, sheep, pigs – 1.0*,**, meat of mammals (except sea mammals) – 0.1*,**, milk – 0.02*,**, poultry meat and offal – 0.01*,**, wheat bran, not processed – 0.3*,**
75	Aminophumare acid dimethyl ester	RNR
76	Amitraz	Cucumbers, tomatoes, fruits (pomaceous and stone types) - 0.5; oranges – 0.5*,**, meat (, pigs, sheep) – 0.2*,**, milk – 0.01*,**, cottonseed (seeds) – 0.5*,** cottonseed (oil, non-refined) – 0.05 – 0.01 ; honey, hop - 0.2
77	Amitrol	Grape, fruits (pomaceous and stone type) – 0.05*,**
78	Arachidonic acid	RNR
79	Atrazine	Corn (grain) -0.03; meat, eggs -0.02;milk –

		0.05
80	Acetoxime	NR
81	Acetamipride	Cereal grain, potatoes -0.5; cucumbers, tomatoes -0.3
82	Poliprenol acetates (made of needles of Siberian fir)	RNR
83	Acetylenic alcohol	RNR
84	Acetochlorine	Soya (beans), sunflower (seeds), rape (grains, oil) -0.01; soya (oil) 0.04; sunflower (oil) -0.02; corn (grains) 0.03
85	Acephate	Artichoke – 0.3*,**; beans -5.0*,**, cabbage – 2.0*,**; cranberry – 0.5*,**; pepper Chili (dry) – 50.0*,**; poultry: fat – 0.1*,**, meat – 0.01*,**, offal – 0.01*,**; mammals: meat – 0.05*,**, milk – 0.02*,**; eggs – 0.01*,**; soya (beans, dry) – 0.3*,**; tomato – 1.0*,**
86	Acifluorfen	Soya (beans , oil) -0.1
87	Anaerobic bacterias activated cultures	RNR
88	Benalaxyl	Grape, melon – 0.3*,**; lettuce (head) – 1.0*,**; onion, potato – 0.2*,**; tomato – 0.2*,**; water melon – 0.1*,**
89	Bendiocarb	Sugar beet, corn (grain) -0.05*
90	Benzoyl formic acid sodium salt	Cottonseed plant (oil), flax (seeds), cereal grain –0.5 NR
91	Benzoilpropetil	NR
92	Benzoic acid	All plant products - NR
93	Benomyl	Cereal grain, rice - 0.5; sugar beet-0.1; sunflower (seeds), potatoes-0.1*; grapes (berries, juice), soya (oil)-0.015; vegetables (except potatoes), fruits (pomaceous fruits, stone fruits)– 0.075; soya (beans) – 0.02
95	Bensulide	NR
95	Bensultape	Potatoes, hop, tomatoes, eggplants -0.04; cereal grain-0.05
96	Bensulphuron-methyl	Rice - 0.02
97	Bentazone	Cereal grain, rice, soya (beans, oil) - 0.1; sorghum, potato – 0.1*,**; beans (dry), peanut – 0.05*,**; bulb onion, flax (seeds) – 0.1*,**; green pea – 0.2*,**; corn (grain) – 0.2 0.1 ; peas (dry); eggs – 0.05*,**; meat of mammals (except sea mammals), milk – 0.05*,**; dry hop - 1.0*
98	Beta-ciflutrine	Fruits (pomaceous fruits), potatoes - 0.2; cabbage, cereal grain, rapeseed (grain, oil)-0.1; pea – 0.2*, sugar beet-0.5

99	Bixafen	Cereal grain – 0.5
100	Binapacryl	Fruits, citrus fruits – NR
101	Bioresmetrin	Cereal grain (wheat), flour – 1.0*,**; bran (not processed) – 5.0 *,**, wheat sprouts – 3.0*,**, tomatoes, cucumbers - 0.4; pepper - 0.01*; fish-0.0015; currant – 0.02*
102	Bispyribac acid	Rice -0.2*
103	Sodium bispyribac	Rice-0.1
104	Bitertanol	Fruits (stone type, except plums) – 1.0*,**, bananas, cucumbers – 0.5*,**, cereal grain, meat of mammals (except sea mammals), milk, mammals' offal – 0.05*,**, pumaceous fruits, plums, except prunes) – 2.0 *,**, eggs, poultry (meat, offal) – 0.01*,**, tomato – 3.0*,**
105	Bifenazate	Cottonseed (seeds) – 0.3*,**, raisin, sweet pepper, fruits (stone type), strawberry – 2.0*,**, vegetables with fruits fit for human consumption, pumpkin type, tomato – 0.5*,**, grapes, pomaceous fruits – 0.7*,**, hop (dry) – 20.0*,**, pepper Chili – 3.0*,**, nuts – 0.2*,**, mammals' meat (except sea mammals), milk fat – 0.05*,**, milk, poultry (meat, offal) – 0.01*,**, mint – 40.0*,**, eggs, offal (mammals) – 0.001*,**, almond in shell – 10.0*,**
106	Biphenthrin	Cottonseed (oil) - 0.015; fruits (pomaceous fruits, except pears) -0.04; pear – 0.5; grapes - 0.2; tomatoes, cucumbers - 0.4; corn (grain) - 0.05 0.04 ; sugar beet – 0.05 0.1* ; corn (oil), sunflower (seeds, oil)-0.02; cabbage-1.0; rapeseed (grain. oil)-0.1; grain cereals (stored) – 0.5; fat, meat C – 0.5*,**, kidney, liver, milk C – 0.05*,**, chicken eggs – 0.01*,**, poultry fat, meat, offal, lemon, orange, potato, grapefruit – 0.05*,**, hop (dry) – 10.0*,**, strawberry – 1.0*,**, wheat bran, non-processed – 2.0*,**, wheat flour – 0.2*,**, wholegrain wheat flour -0.5*,**,
107	Boskalide	Pomaceous fruits – 2.0; vegetables with edible roots and tubers – 2.0*,**, bananas – 0.6*,**, cereal grain: barley, oat, wheat, rye – 0.5*,**, other grain crops – 0.1*,**, berries and other small fruits, except strawberries and grape), prunes, pepper Chili (dry), raisin – 10.0*,**, cabbage (all types), vegetables with edible bulbs, kiwi – 5.0*,**, grape – 5.0; coffee

		beans, tree nuts (except pistachio and almond) – 0.05*,**; almond in shell – 15.0*,**; leaf vegetables (lettuce, spinach, parsley, etc.) – 30.0*,**; fruit-bearing vegetables, pumpkin, legumes (beans, peas), fruits (stone type), except prunes, strawberry – 3.0*,**; mammals' meat (except sea mammals) – 0.7*,**; mammal's offal – 0.2*,**; eggs, poultry meat, fat, offal – 0.02*,**; milk – 0.1*,**; milk fat – 2.0*,**; pistachios – 1.0*,**; oilseeds – 1.0*,**; sunflower (seeds), rapeseeds (seeds) – 1.0*,**; sunflowerseed (oil)-0.5; rapeseed (seed, oil)-0.2
108	Brodifacum	RNR
109	Bromadiolone	RNR
110	Bromide 4- methyl benzole aldehyde triphenyl-phosphonium +4-nitrodiphenylazo-metin methylenetriphenyl- phosphonium - bromide	Corn NR
111	Bromoxynil	Cereal grain, millet, corn (grain) -0.05
112	Bromophos	Cabbage, frigole, cucumbers, salad, pea, grapes -0.05; fruits (pomaceous fruits) -0.1; fruits (stone fruits)-0.07; dry hop - 0.5; berries- 0.04
113	Brompropilate	Grapes – 2.0* 0.01* ; citrus fruits, pomaceous and stone fruits 2.0 0.03, 0.02 ; pulses(pods and seeds, not ripe) – 3.0*,**; cucumbers, pumpkin, melon – 0.5*,**; fruits (stone type, except prunes), strawberry – 2.0*,**; berries – 0.05; cottonseed plant (oil) -0.02*; honey - 0.02
114	Bromuconazol	Cereal grain, fruits (pomaceous fruits), grapes - 0.04; berries - 0.08
115	Bronopol	NR
116	Bupirimat	Cucumbers, melons, fruits (pomaceous fruits)- 0.1 currant-0.1
117	Buprofezin	Almond – 0.05*,**; almond in shell – 2.0*,**; pomaceous fruits: apples – 3.0*,**, pears – 6.0*,**; stone type fruits: cherry, plums (except prunes) – 2.0*,**, peache, nectarine – 9.0*,**; citrus fruits, grape – 1.0*,**; tomato – 1.0 0.2 ; strawberry – 3.0*,**; dried pulp of citrus fruits, raisin, pepper – 2.0*,**; meat and offal of mammals (except sea) – 0.05*,**; pumpkin – 0.7*,**; cucumbers – 0.7 0.1 ; mango – 0.1*,**; milk – 0.01*,**; olives –

		5.0*,**, pepper Chili (incl. dry) – 10.0*,**
118	Butylate	Corn (grain) -0.5*
119	Butoxicarboxim	Citrus fruits (pulp) - 0.01
120	Vamidothion	Vegetables (except for potatoes) -0.2
121	Vernolat	Soya (beans), corn(grain) -0.5*; soya (oil) - 0.1*; tobacco- 1.0*
122	Vinclozolin	Blackberry – 5.0*,**; cabbage – 1.0*,**; cattle meat and milk – 0.05*,**; cauliflower – 1.0*,**; cherry – 5.0*,**; chicken eggs – 0.05*,**; chicory - 5.0*,**; pulses – 2.0*,**; cucumbers – 1.0; currant (red, black, white) – 5.0*,**; orchard peas – 1.0*,**; gherkin – 1.0*,**; gooseberry – 5.0*,**; grape – 5.0 3.0; hop (dry) – 40.0*,**; kiwi – 10.0*,**; lettuce – 5.0*,**; melon – 1.0*,**; pepper Chili – 1.0*,**, sweet pepper – 3.0*; pomaceous fruits – 1.0*,**; potato – 0.1*,**, rapeseed (seeds) – 1.0*,**; raspberry (black, red) – 5.0*,**; strawberry – 10.0*,**; tomatoes – 3.0 1.0*; sunflower (seeds and oil) -0.5*
123	Granulovirus admixed with polyhedrosis of turnip moth	RNR
124	Granulovirus of apple worm	RNR
125	Nuclear polyhedrosis virus of cabbage moth	RNR
126	Nuclear polyhedrosis virus of lackey moth	RNR
127	Nuclear polyhedrosis virus of gypsy moth	RNR
128	Nuclear polyhedrosis virus of cotton budworm	RNR
129	Hydrogen phosphide	Cocoa beans, dry fruits and vegetables, peanuts, spices, tree nuts – 0.01*,**; cereal grain – 0.1*,**
130	Galaxifop	Banana, coffee beans, stone type fruits – 0.02*,**; citrus fruits, grape, pomaceous fruits – 0.05*,**; onion, bulb – 0.2*,**
131	Galaxifop-P methyl	Sugar beet, sunflower (seeds), soya (beans), vegetable oil -0.05; cottonseed seeds – 0.05*,**; rapeseed (grain) - 0.2; potatoes- 0.01
132	Galaxifopetoxiethyl	Sugar beet, sunflower (seeds), soya (beans), vegetable oil -0.05; cottonseed (seeds) -0.05*; rapeseed (seeds) - 0.2; potatoes - 0.01*
133	Gamma- Cyhalothrin	Cereal grain -0.05; rapeseed (grain, oil), fruits (pomaceous fruits)-0.1; potatoes, carrot, sugar beet -0.02; onion – 0.2

134	Hexaflumuron	Potatoes - 0.05
135	Hexachlorbenzene	Cereal grain -0.01
136	Hexachlorobutadiene	Grapes and products of its processing – 0.0001
137	Hexachlorocyclohexane (α , β , γ -isomers) (HCCH)	Meat and poultry (fresh, cooled, and frozen) - 0.1; byproducts (liver, kidneys) -0.1; sausages, cookery, Meat and poultry cans - as per raw materials (in terms of fat); eggs, gelatin -0.1; milk and fermented milk products -0.05; milk processing products (cheeses, curd products, butter, cream, sour milk), concentrates of milk and whey proteins, milk and dry milk products (in terms of fat) - 1.25; fresh water fish (fresh, cooled, and frozen) -0.03; sea tuna fish (fresh, cooled, and frozen), meat of sea animals - 0.2; salty, smoked, air-dried fish -0.2; fish cans (fresh water, sea, tuna fishes, meat of sea animals) - as per raw materials; liver of fishes and its derived products, cans fish liver - 1.0; caviar, fat herring - 0.2; cereal grain, pulses - 0.5; flour, grits -as per raw materials; soya, corn (grain), bakery confectionery products – 0.2; starch and syrup made of corn-0.5; starch and syrup made of potatoes, sugar beets -0.1; flax (seeds), rape (grain), mustard - 0.4; sunflower (seeds), peanut, nuts, cocoa (beans), cocoa-products - 0.5; non-deodorized oil - 0.2; deodorized oil, of best purification - 0.05; animal fat - 0.2; fish fat-0.1; vegetables, melons and gourds, mushrooms - 0.5; potatoes - 0.1; fruits, berries, grapes - 0.05; cans with fruits and vegetables - as per raw materials; juices - as per raw materials; honey -0.005; protein products of seed corn, grain legumes seeds, and seeds of other crops-0.1; baby products: adapted milk mixes for children 0 - 3 months) -0.02; products for children 4 - 12 months: milk – 0.02; cottage cheese 18% - 0.1; meat - 0.02; grits, vegetables, potatoes, fruits - 0.01; butter - 0.2; vegetable oil- 0.01
138	Hexithiatox	Citrus fruits – 0.5* (pulp – 0.02*); cottonseed (oil) - 0.1 *; fruits (pomaceous fruits) – 0.4*, grapes - 0.1*; strawberry – 0.5*, **; dates, hop (dry) – 2.0*, **; raisin, prunes -1.0*, **; mammals offal, eggs, mammals fat (including milk fat), milk, meat of mammals (except sea mammals), poultry meat and offal, vegetable

		with edible fruits, pumpkin type, except water melons, tree nuts – 0.05*,**; eggplants, tomato – 0.1*,**; grape meal (dry) – 15.0*,**, stone type fruits – 0.3*,**
139	Heptachlor	Cereal grain – 0.02*,**; citrus – 0.01*,**; cottonseed (seed) – 0.02*,**; eggs – 0.05*,**; meat of mammals (except sea mammals) – 0.02*,**; milk – 0.006*,**; pineapple – 0.01*,**; poultry meat – 0.02*,**; soya (beans) – 0.02*,**; soybean oil, crude – 0.05*,**, soybean oil, refined – 0.02*,** All food products – 0.007
140	Gibberellic acid sodium salts	RNR
141	Gibberellin -A 3	RNR
128	Maleic hydrazide	potatoes – 20; onion – 15; sugar beet, red beet, garlic, carrot, tomatoes, water melons – 8.0; green tobacco – 30
142	Hymexazol	Sugar beet, red beet - 0.01
143	Glyphosate	Fruit (pomaceous fruit, stone fruit), citrus fruit, sunflower (seeds) , vegetables, potato, corn (grain), mushrooms- 0.3; watermelon - 0.3*; grapes, berries (including wild berries) -0.1; cereal grain- 30.0 3.0 ; rice -0.15*; bananas – 0.05*,**; corn (grain), soya (dry beans) – 5.0; sunflowerseed (seeds) – 7.0; rapeseed (seed) – 20.0; sunflower oil, rapeseed oil, soybean oil – NR; peas (dry) – 5.0; cottonseed (seed) – 40.0*,**; mammals' offal – 5.0*,**; eggs, meat of mammals (except for sea mammals), poultry meat, milk – 0.05*,**; pig offal and poultry offal – 0.5*,**; beans (dry), sugar cane – 2.0*,**; sugar cane molasses – 10.0*,**; wheat bran, not processed – 20.0*,** soybean oil-0.05*; – sunflower seed oil-0.1; soya bean-0.15
144	Glyphosate trimesium	Cereal grain, fruit (pomaceous fruit), grapes - 0.3
145	Glufosinate ammonium	Fruit (seeded fruit, stone fruit), berries and other small fruits (except currant), citrus fruits, grapes, carrot - 0.2; potatoes – 0.5 0.2 ; sunflower (seeds), rapeseed (seeds) – 5.0; buckwheat, millet, cereal grain – 0.4; vegetable oil, except crude sunflower and rapeseed oil) – 0.4; pea – 3.0, soya (beans), beans – 2.0; almond (in shell), currant (black, red, white) – 0.5*,**; asparagus, tropical and sub-tropical

		fruits, except bananas, beans, corn – salad, eggs, meat of mammals (except sea mammals), onion (bulb), sugar beet, poultry meat , non-refined rapeseed and sunflowerseed oils – 0.05*,**; bananas – 0.2*,**; edible offal of mammals and poultry, corn, tree nuts – 0.1*,**, milk – 0.02*,** sunflower (seeds); rapeseed (seeds); rapeseed, leguminous – 0.4;
146	Guazatine	Cereal grain - 0.05; citrus fruits – 5.0*,**
147	Humic acids	RNR
148	Ammonium salt of humic acid	RNR
149	Sodium salts of humic acids	RNR
150	A (+) - (p-nitrophenyl) - 1 ,3-dihydroxy isopropyl-ammonium-2-chlorethylphosphoric acid	Tomatoes- 1.5
151	DAER	Grapes, sugar beet - 0.1; red beet, cottonseed oil - 0.5; citrus (pulp)- 0.05
152	Dazomet	Potato, vegetables, fish - 0.5
153	Dalapon	Fruit (seeded fruit, stone fruit), grapes, potatoes, red beet, sugar beet -1.0; cottonseed oil - 0.1; tea-0.2; berries (including wild) – 0.6
154	Daminozide	Fruit (seeded fruit) -3.0
155	Deltametrin	Sunflower (seeds), melon, tobacco-0.1*; cottonseed oil, sunflower seed oil, bananas - 0.05*; fruit (stone fruit) – 0.2*,** 0.05* cereal grain – 0.2, berries (except strawberry) – 0.5**; strawberry – 0.2**; leguminous, beans (dry) – 1.0, fruit (seeded fruit) , cabbage, corn (grain), cucumbers, lettuce, rice, citrus fruit (pulp) , sugar beet – 0.01; potatoes, tomatoes, grapes, carrot – 0.01 ; watermelon, soya-bean oil, pepper, cacao beans, -0.01*; potato – 0.1; dry hop -5.0*; meat, milk 0.02 ; liver, kidneys (cattle, goat, pig, sheep), milk - 0.05; meat (except mammals and poultry) , rapeseed (seed and oil), corn (oil), citrus fruits, carrot – 0.02; animal fat- 0.5; tomato – 0.3, bean-type vegetables with edible fruits, pumpkin type, leek – 0.2*,**; eggs, poultry offal, hazel-nut, sweet corn (boiled and in cobs), walnut – 0.02*,**; broccoli, Chinese and cauliflower cabbage – 0.1*,**, leaf-type vegetables, wheat flour, non-screened – 2.0*,**; lentils (dry), olives – 1.0*,**; meat (mammals), except sea animals – 5.0*,**; mushrooms, onion (bulb) –

		0.05*,**; poultry meat -0.1*,**; radish 0.01*,**; tea (green and black), wheat bran (not processed) – 5.0*,**; wheat flour – 0.3*,**; pomaceous fruits, grape – 0.2
156	Demeton	Cereal grain, cottonseed oil -0.35
157	Desmedipham	Red beet, sugar beet - 0. 1
158	Desmetryne	Cabbage - 0.05; onion - 0.05*
159	Diazinon	Cereal grain, onion, potatoes, cottonseed oil, corn (grain), rutabaga, turnip, red beet, sugar beet - 0.1; cabbage 0.1 , tobacco, cucumbers, tomatoes, poppy seed oil -0.5; dry hop - 1.0; walnut – 0.01*,**; almond, sweet pepper, Chinese cabbage, pumpkin – 0.05*,**; blackberry, dewberry, strawberry, pineapple, radish – 0.1*,**, musk melon, raspberry, currant (red, black, white), cranberry, peach, kiwi, kohlrabi, peas (fresh beans), beans (pods/seeds) – 0.2*,**; pomaceous fruits – 0.3*,**; pepper Chili,(dry), broccoli, lettuce and leaf salad, spinach – 0.5*,**; pineapple, strawberry, plums (except prunes), cherry, onion –shallot – 1.0*,**; prunes – 2.0*,**; eggs and poultry meat – 0.02; sweet corn (boiled, in cobs),poultry offal – 0.02*,**; meat of C, goat, pigs, sheet – 2.0; liver and kidney of C, goat, pigs, sheep – 0.03*,**; milk (dairy products) – 0.02 carrot -0.08; meat (in terms of fat), milk, milk products, poultry, eggs—0.01
160	Diafentiuron	Cucumbers, tomatoes -0.05;
161	Dibromo-chloro propane	NR
162	Potassium salt of diisopropyldithiophosphonic acids (1-Hydroxyethylidenediphosphonic acid)	Cereal grain —NR
163	Dicamba	Cereal grain, corn (grain)-0.5; corn oil - 0.05;
164	2-ethylhexyl ether of dicamba	millet -0.3 NR
165	Diquat (dibromide)	Pea – 0.2, carrot, potatoes - 0.05; sunflower (seeds) – 1.0, rape (seeds)- 2.0 0.5 ; sunflower seed oil, rapeseed oil, soya-bean oil -0.1; soya bean 0.2; milk – 0.01* 0.4 ; barley – 5.0*,**; beans, lentils (dry), rice, milled – 0.2*,**; meat of mammals (except sea mammals), mammals' offal, eggs, corn, poultry meat and offal, vegetable oil , rude (except sunflowerseed, soybean and rapeseed oil), vegetables with edible tubes, roots, fruits and bulbs – 0.05*,**; rice – 10.0*,**, rice milled – 1.0*,**; wheat

		bran not processed, wheat flour, not sifted, wheat, oat, sorghum – 2.0*,**; wheat flour – 0.5*,** meat – 0.01
166	Dichloran	Fruit (stone fruit) (peach, nectarine) – 7.0 0.1* ; fruit (seeded fruit) – 0.06; carrot – 15.0, onion (bulb) – 0.2; cabbage, potatoes – 0.004*, grape – 7.0*
167	Diclofop methyl	Sugar beet -0.01; soya bean -0.05; soya-bean oil - 0.02*
168	Dicofol	Pepper – 1.0, tomatoes – 0.1*, cucumbers – 0.5, fruit (seeded fruit) – 0.1*, fruit (stone fruit): peach, cherry – 5.0, plums – 1.0, grapes – 5.0, eggplant – 0.1*; pumpkin – 1.0, citrus fruit (pulp) – 5.0 0.1* ; dry hop – 50 5.0 ; berries – 0.05; cottonseed oil – 0.5 0.01* ; beans (dry beans) – 0.1*,**; beans (pods and/or seeds) – 2.0*,**; melon – 0.2*; pepper Chili (dry) – 10.0*,**; prunes – 3.0*,**; cottonseed (seed) – 0.1*,**; cottonseed oil, non-refined – 0.5**, walnut, pecan – 0.01*,**; milk – 0.1*,**; eggs – 0.05*,**; meat (cattle) - 3.0*,**; offal (cattle) – 1.0*,**; poultry meat (0.1*,**); poultry offal – 0.05*,**; tea (green and black, fermented, dried) – 50.0*,**
169	Dimethylchlor	Rape (seeds, oil) -0.02*
170	Dimethenamid -P	Corn (grain), soya bean, soya-bean oil, sugar beet, red beet, beans (dry) -0.02; sunflower (seeds, oil) -0.04; potato, garlic, onion (bulb), onion shallot, sorghum, sweet corn (boiled cobs), sweet potato, peanuts, eggs, meat of mammals (except sea mammals), milk, poultry neat and offal – 0.01*,**
171	Dimetipin	Sunflower (seeds) – 1.0; sunflower seed oil - 0.05*; potatoes – 0.05; rapeseed (seed) – 0.2*,**; cottonseed seed – 1.0*,**; cottonseed oil, non-refined, cottonseed oil (for human consumption) – 0.1*,**; meat of mammals (except sea mammals), poultry meat, offal, eggs, milk – 0.01*,**
172	Potassium salt of dimethyl ether of dehydro-aspartic acid	NR
173	Dimethoate	Artichoke – 0.05*,**; asparagus – 0.05*,**; barley – 2.0*,**; Brussels cabbage – 0.2; savoy cabbage – 0.05*,**; cauliflower – 0.2*,**; CATTLE offal – 0.05*,**; celery – 0.5*,**; cherry – 2.0; citrus fruits – 5.0; eggs –

		0.05*,**; salad – 0.3*,**; CATTLE fat, except milk fat – 1.0*,**; mango – 1.0*,**, meat of cattle, horses, pigs, goats, sheep – 0.05*,**, olives – 0.5*,**, pears – 1.0; peas, beans – 1.0*,**; pepper Chili – 3.0*,**; sweet pepper, including pimento, - 0.5*,**, potato – 0.05; poultry fat – 0.05*,**, chicken offal – 0.05*,**, sheep offal – 0.05*,**, sugar beet – 0.05; wheat – 0.05*, olives, mushrooms, rice, melons type, red-beet, cucumbers, tomato, tobacco, dry hop, berries, legumes, millet, grape, sunflowerseed (seeds, oil) – 0.02; rapeseed (seeds, oil) – 0.05 fruit (seeded fruit, stone fruit), olive, mushrooms, rice, gourds, cucumbers, tomatoes, tobacco, sugar beet, red beet, dry hop, berries, cabbage, cereal grain, leguminous, millet, grapes, citrus fruits, potatoes, sunflower (seeds), sunflower seed oil -0.02; rape (seeds, oil) – 0.05
174	Dimetomorf	Broccoli – 1.0*,**; cabbage – 2.0*,**; corn salad – 10.0*,**; grapes – 3.0; raisin – 5.0*,**, mammals' offal – 0.01*,**, eggs – 0.01*,**, fruit bearing vegetables, except pumpkin – 1.0*; pumpkin – 0.5*,**, cucumbers – 0.01. dry hop – 80.0*,**, kohlrabi – 0.02*,**, salad – 10.0*,**, meat of mammals (except sea mammals) – 0.01*,**, milk – 0.01*,**, pepper Chili (dry) – 5.0*,**, pine apple – 0.01*,**, potato – 0.5; poultry meat and offal – 0.01*,**, strawberry – 0.05*,** potatoes – 0.5; cucumbers – 0.01; grapes – 3.0
175	Dimoxystrobin	Sunflower (seeds), sunflower seed oil, rape (seeds, oil) -0.05
176	Diniconazole	Cereal grain - 0.05
177	Dinitroorthokrezol	Cucumbers, potatoes, grapes - 0.06; dog rose – 0.1
178	Dinobuton	Tomatoes, cucumbers, fruit (seeded fruit), grapes, sugar beet citrus fruit (pulp), cottonseed oil, pepper -0.05; berries -0.05; dry hop - 0.5
179	Dinokap	Cucumbers – 1.0; vegetables with edible fruits, pumpkin type – 0.05*, fruit (seeded fruit – 0.2*; grapes – 0.5* 1.0 ; strawberry – 0.5; peach – 0.1*,**; pepper – 0.2*; pepper Chili (dry) – 2.0*; tomato – 0.3*,** berries – 0.2
180	Dipropetrin	Watermelon -0.1

181	Disulfoton	Cereal grain: barley, wheat – 0.2*,**; oat – 0.02*,**; beans (in pods and/or non-ripened), beans (dry) – 0.2*,**; corn (grain), sweet corn (boiled cobs), sweet corn (grain) – 0.02*,**, sugar beet – 0.2*,**, peas (pods, non-ripened seeds) – 0.1*,**, peas green, juicy seeds) – 0.02*,**, nuts (peanut, pecan) – 0.1*,**, pine apple – 0.1*,*, coffee beans – 0.2*,**, cottonseed seed 0.1*, asparagus – 0.02*,**, poultry meat – 0.02*,**, milk (cattle, goat, sheep) – 0.01*,**
182	Ditalimfos	Cereal grain, cucumbers -0.1; fruit (seeded fruit), grapes -0.5; berries – 0.02
183	Dithianon	Fruit (stone fruit) (cherry and other)- 5.0* 0.02* ; grapes -3.0 1.5 ; citrus fruits (mandarins, grapefruit, pommels, etc.) - 3.0*,**; berries, small fruits – 5.0*,**; fruit (seed fruit) – 5.0* 2.0
184	Dithiocarbamates	Nuts (almond, pecan), peanuts, asparagus - 0.1*,**; almond in shell – 20.0*,**; bananas, cucumbers, mango, oranges, tomato – 2.0*,**; cereal grain, carrots, sweet pepper, pumpkin (early), water melon – 1.0*,**; cabbage, cranberry, papaya fruits (seed type), strawberry – 5.0*,**; cherry, potato, pumpkin – 0.2*,**; salad, currant (red, black, white), mandarins, pepper Chili (dry) -10.0*,**; garlic, leek, salad, melon (except water melon), onion, shallot 0.5*,**; leaf cabbage – 15.0 *,**, hop (dry) – 30.0*,**; fruits (stone type), except cherry – 7.0*,**, sweet corn (in cobs) -0.1*,**, meat of mammals (except sea mammals), milk, eggs – 0.05*,**; offal of mammals, poultry meat and offal – 0.1*,**
185	Diuron	All food products – 0.02
186	Diphenamid	Tomatoes, pepper - 0.1; tobacco -0.15;
187	Diphenylamine	Seed type fruits: apples -10.0*,**, pears – 5.0*,**; apple juice – 0.5*,**; meat, kidney (cattle) – 0.01*,**; liver (cattle) – 0.05*,**; milk, milk fat – 0.01*,**
188	Difenoconazole	Fruit (seeded fruit) – 1.0, sugar beet, red beet - 0.2; cereal grain -0.08; fruit (stone fruit), except nectarines and peaches) – 0.2; nectarines and peaches – 0.5- 0.15 ; tomatoes – 0.6 0.05 ; carrot -0.3; potatoes -0.02; celery – 5.0**, grape – 0.5; asparagus – 0.03*,**;

		bananas – 0.5**; citrus – 0.6**; rice – 1.0**; broccoli – 0.5*,**; cabbage (Brussels, cauliflower, cabbage), mammals offal, papaya – 0.2*,**; mango – 0.07*,**; eggs, poultry meat and offal – 0.01*,**; garlic, soya (beans), sunflowerseeds (seeds) – 0.02*,**; leek – 0.3*,**; salad leaf and head, olives – 2.0*,**; meat of mammals (except sea mammals), rapeseed (seeds) – 0.05*,** , milk – 0.005*,**
189	Diflubenzuron	Seed type fruit – 5.0; mushrooms (incl. champignon) - 3.0; cabbage – 1.0; citrus fruit – 0.5*,**; meal and offal of mammals (except sea mammals) – 0.1*,**; eggs, poultry meat – 0.05*,**; milk – 0.02*,**; rice – 0.01*,** fruit (seeded fruit) champignon – 0.1; cabbage – 1.0
190	Diflyufenikan	Cereal grain – 0.05
191	Diclobutrazol	Cereal grain -0.1*
192	Dichloral urea	NR
193	Dichlorprop dichlorprop-P	Cereal grain, flour - 0.05
194	Dichlorphos	Cereal grain – 5.0; wheat bran – 10.0; cabbage, fruit (seeded fruit, stone fruit), citrus fruit (pulp), grapes, berries, tea -0.05; cereal groats, livestock products – 0.1*,** wheat flour – 1.0*,** , wheat sprouts – 10.0*,**; coarse- milled flour – 2.0*,** grain, bran – 0.3; livestock products, cereals – 0.01
195	Dichlofluanid	Seed type fruits – 5.0; berries: currant (red, black, white) raspberry – 15.0; strawberry – 10.0; gooseberry – 7.0; grapes – 15.0; cucumbers – 5.0*,**; lettuce – 10.0*,**; onion (bulb) – 0.1*,**; potato – 0.1*,**; tomato – 2.0*,**; peach – 5.0*,**; pepper – 2.0*,**; pepper Chili (dry) – 20.0*,** berries, grapes, fruit (seeded fruit) – 0.02
196	Dichloropropene + dichloropropane	NR
197	Dicyandiamide (metabolite and half- product of synthesis of Granstar)	NR
198	Dodin	Cherry – 3.0*,**; nectarine – 5.0*,**; peach – 5.0*,**; pomaceous fruits – 5.0*,**
199	Doramectin	For cattle: meat -0.01;fat-0.15; liver-0.1; kidneys-0.03; for sheep and pigs: meat - 0.01;fat-0.1; liver-0.05; kidneys -0.03
200	Zoxamide	Dried grape (raisin, kishmish, etc.) – 15.0*,**; vegetables with eatable fruits, pumpkin type – 2.0*,**; grapes – 5.0*,**; potato – 0.02*,**; tomato – 2.0*,**

201	Ivermektin	For cattle: fat-0.04; liver-0.1; meat - RNR; for sheep and pigs: fat-0.02; liver-0.015; meat-RNR; meat and offal of poultry-0.001
202	Isobutene dichlorides (mixture)	NR
203	Isoxadifen-ethyl	Corn (grain and oil) - 0.2
204	Isoxaflutole	Corn (grain) - 0.05; corn (oil) – 0.1
205	Isopropalin	Tobacco - 1.0*
206	Izopropilfenatsin	RNR
207	Izoprotiolan	Rice - 0.3
208	Isoproturon	Cereal grain -0.01
209	Isofenphos	Rapeseed - NR
210	Imazakvin	Soya bean, soybean oil - 0.1*
211	Imazalil	Banana- 2.0*,**; citrus fruits – 5.0*,8*;; cucumbers (incl. gherkins) – 5.0*,**, melon – 2.0*,**; persimmon Japanese – 2.0*,**; pomaceous fruits – 5.0*,**; berries (raspberry (red and black), strawberry, and other – 2.0*,**; cereal grain (wheat and other) – 0.1; soya bean - - sunflower (seeds), rape (seeds) – 0.02; soya-bean oil, sunflower seed oil, rapeseed oil -0.04; corn (grain)-0.3; millet – 0.4
212	Imazametabenz	Cereal grain -0.2
213	Imazamox	Soya bean, soybean oil, pea -0.05; rape (seeds, oil) -0.1; sunflower (seeds and oil)-0.1
214	Imazapyr	Wild berries -2.0; wild mushrooms-4.0; sunflowerseeds (seeds, oil) – 0.1
215	Imazetapir	Soya (beans, oil), peas – 0.5
216	Imidaclopride	Almonds (in shell) – 5.0*,**; fruits (pomaceous fruits), except pears-0.5; pears – 1.0; apple meal, dry – 5.0*,**; stone type fruits (peaches, cherry, nectarine, apricot) – 5.0*,**, plums (including prunes) – 0.2*,**; bananas – 0.05*,**; beans – 2.0*,**, berries and other small fruits (orchard strawberry, cranberry, other) – 3.0; cabbage (all types)-0.5; cereal grain -0.1; citrus fruits – 1.0**; citrus fruits (dry pulp) – 10.0*,**; coffee (beans) – 1.0*,**; cucumbers-1.0; offal of mammals -0.3*,**; egg-plants-0.5** eggs – 0.02*,**; grape – 1.0; hop (dry) -10.0*; onion (bulb, green, leek) – 0.2; salad – 2.0*,**; mango – 0.2*,**; meat of mammals (except sea mammals) – 0.1*,**; melon – 0.2*,**; milk – 0.1*,**; peanuts – 1.0*,**; peas (dry, shelled, sweet, raw pods, seeds) – 5.0; nuts (pecan) – 0.05*,**; pepper –

		1.0**. Pepper Chili (dry) – 10.0**, **; pomegranate – 1.0**, **; poultry meat – 0.05**, **; rape (grain, oil) -0.1, vegetables with edible roots and tubers – 0.5**, **; squash – 1.0*; sunflower (seeds)-0.4; sunflower (oil)- 0.2; soybeans (seeds, oil) – 0.1; corn (sweet (boiled in cobs) – 0.02**, **; tomatoes – 0.5; water melon – 0.2**, **; wheat bran, not processed – 0.3*; wheat flour – 0.03**, **; carrot, red beet, sugar beet, , tomatoes, potatoes – 0.5; corn (grain, oil) – 0.1; black currant 3.0; berries 3.0**, **; pepper 1.0**, **;
217	Indoxacarb	Apples – 0.5; broccoli – 0.2**, **; cabbage – 3.0**, **; cauliflower – 0.2**, **; cranberry – 1.0**, **; raisin – 5.0**, **; offal of mammals, edible – 0.05**, **; eggplant – 0.5**, **; eggs – 0.02**, **; pumpkin – 0.5**, **; grapes – 2.0*, salad (head) – 7.0**, **; salad leaf – 15.0**, **; meat of mammals (except sea mammals) – 2.0**, **; milk fat – 2.0**, **; milk 0.1**, **; mint – 15.0**, **; ground nut – 0.02**, **; pear – 0.2*; pepper – 0.3**, **; potato – 0.02**, **; poultry meat and offal – 0.01**, **; prunes – 3.0**, **; soybeans (beans, dry) – 0.5**, **; tomato – 0.5; rapeseed (seeds, oil) – 0.05; onion – 2.0 Fruits (seeded fruits), grapes 0.5
218	Iodfenfos	Cabbage, gooseberries, grapes - 0.5; berries – 0.01
219	Ioxinil	Garlic, onion -0.1
220	ipkonazole	Cereal grain-0.02
221	Iprobenfos	NR
222	Iprodione	Almonds – 0.2**, **; barley – 2.0**, **; beans (dry) – 0.1**, **; berries (blackberry, strawberry, raspberry black and red) – 15.0**, **, cabbage (broccoli, Chinese and other) – 5.0**, **; carrots – 0.5; stone type fruits: cherry, peach and other – 10.0**, **; seed type fruits – 5.0**, **; beans whole (pods and seeds) – 2.0**, **; cucumbers – 2.0; grapes – 10.0; kiwi – 5.0**, **; salad (head and leaf) – 10.0**, **; onion (bulb) – 0.2**, **; sugar beet – 0.1**, **; tomatoes – 5.0; chicory sprouts – 1.0**, **; rapeseed (seeds) – 0.5**, **; rice, milled – 10.0**, **; sunflowerseed (seeds) – 0.5; sunflowerseed (oil) – 0.02; potato – 0.05 grapes – 0.4; cucumbers, sunflower (seeds, oil) –

		0.02; potatoes, carrot—0.05; tomatoes—5.0; eelery cabbage—5.0**; lettuce—10.0** berries—15.0**
223	Isazofos	Tomatoes, cucumbers, berries-0.2
224	Iodosulfuron-methyl-sodium	Cereal grain - 0.1; corn (grain and oil) -0.2
225	Cadusafos	Bananas – 0.01*,**; potato – 0.02*,**
226	Potassium vinyloxy- ethyl dithyocarbamate	Cucumbers - 0.1
227	Captan	Almond – 0.3*,**; blackberry, huckleberry – 20.0*,**; cherry – 25.0*,**; cucumbers – 3.0*,**; dry grapes (all types) – 50.0*,**; grapes – 25.0*; water melon – 10.0*,**; nectarine – 3.0*,**; peach – 20.0*,**; plums – 10.0*,**; seed type fruits – 3.0 (K); potato – 0.05*,**; raspberry – 20.0*,**; strawberry - 15.0*,**; apple juice- 0.01; grapes, grape juice – 0.05 ; fruit (seeded fruit) 3.0
228	Carbaryl	Almonds (in shell) – 50.0*,**; asparagus, citrus fruits – 15.0*,**; red beet, corn (oil, crude), corn (sweet, in cobs) – 0.1*,**, carrot, chili pepper – 0.5*,**; cranberry, sweet pepper (including in pods), tomato – 0.5*,**; eggplant, tree nuts, turnip – 1.0*,**; sweet potato – 0.02*,**; rice: polished – 1.0*,**; rice in shell – 50.0*,**, rice not milled – 170.0*,**; meat of mammals (except sea mammals), milk – 0.05; dairy products – 0.02; kidney of mammals – 3.0*,**; liver of mammals 1.0*,**; olive oil (refined) – 25.0*,**; olives – 30.0*,**; Chili pepper (dry) – 2.0*; sorghum, tomato paste 10.0*,**; soybeans (beans and oil crude), sunflowerseed (seeds) – 0.2*,**; sunflowerseed oil, crude – 0.05*,**; soybeans in shell – 0.3*,**; tomato juice – 3.0*,**; cereal grain (wheat), bran, not processed (wheat) – 2.0*,**; wheat flour – 0.2*,**; wheat sprouts – 1.0*,**; cottonseed (oil) – 0.0125; corn – 0.02; pomaceous fruits, potato – 0.05 cottonseed oil, corn (grain)–0.0125; fruit (seeded fruit, potatoes–0.05; meat 0.01; milk and milk products–0.02
229	Carbendazim	Sugar beet- 0.1; cereal grain - 0.5; strawberry, currant – 1.0; pomaceous fruits – 0.2*,**; grape – 3.0; cucumbers – 0.05*; apricots, nectarines, peaches, Chili pepper, shelled rice – 2.0*,**; asparagus, bananas, carrot – 0.2*,**;

		beans (dry), Brussels cabbage, beans (ordinary, in pods and/or not ripened), plums (including prunes), soya (beans), pumpkin (ordinary), tomato – 0.5*,**; berries and other small fruits (except grapes), oranges (including hybrids) – 1.0*,**; meat of cattle and poultry, chicken fat, offal of mammals, eggs, gherkin, rapeseed (seed), milk – 0.05*,**; cherry – 10.0*,**; coffee beans, peanuts, tree nuts – 0.1*,**; peas orchard, shelled – 0.02*,**;; salad (head), mango, pineapple – 5.0*,**; Chili pepper (dry) – 20.0*,**; cereal grain – 0.2; strawberry, currant – 0.05; fruit (seeded fruit) – 0.05; grapes, cucumbers – 0.05*
230	Carboxin	Corn (grain), millet, cereal grain, potatoes - 0.2; corn oil – RNR
231	Carbosulfan	Potato – 0.25; sugar beet – 0.3; corn – 0.05; citrus, citrus pulp (dry) – 0.1*,**; cottonseed (seed) – 0.05*,**; meat of mammals (except sea mammals), offal of mammal, Poultry meat, eggs and offal – 0.05*,** (control on Carbosulfan and its metabolites) corn (grain), sugar beet – 0.05; potatoes – 0.25 (check of carbosulfan and its metabolites)
232	Carbofuran	Sugar beet – 0.2; rapeseed (seed, oil) – 0.05; dry hop – 5.0*; banana – 0.1*,**; citrus fruits – 0.5*,**; pulp of citrus fruits (dry) – 2.0*,**; corn – 0.05*; coffee beans – 1.0*,**; sugar cane, cottonseed (seed), sorghum – 0.1*,**; sunflowerseed (seed) – 0.1*,**; rice milled (0.1*,**; meat, fat and offal of Cattle, goat, horses, pigs, sheep 0.05*,**; sugar beet – 0.05; dry hop – 5.0*; rape (seeds, oil) – 0.1; mustard (seeds, oil) – 0.05
233	Carfentrazone-ethyl	Cereal grain, rape (seeds, oil), sunflower (seeds and oil), corn (grain and oil) -0.02
234	Quizalofop-P-tefuryl	Potatoes, carrot, tomatoes, cabbage, sunflower (seeds), soya bean, sugar beet, red beet - 0.04; onion, sunflower seed oil, soya-bean oil -0.06; rape (seeds, oil) –0.02
235	Quinmerac	Rapeseed (seed, oil) – 0.1
236	Quinclorac	Rice –0.05
237	Quinoxifen	Barley; wheat – 0.01*,**; cherry – 0.4*,**; strawberry, black currant, dry hop, pepper – 1.0*,**; grape – 2.0*,**; salad (head) – 8.0*,**; salad (leaf) – 20.0*,**; melon –

		0.1*,**; pepper Chili (dry) – 10.0*,**; sugar beet – 0.03*,**; offal of mammals and poultry, milk, eggs – 0.01*; meat of mammals (except sea mammals), milk fat – 0.2*,**; poultry meat – 0.02*,**
238	Quintozene	Barley, soya beans, cottonseed seed, corn, peas, dry sugar beet – 0.01*,**; broccoli, sweet pepper – 0.05*,**; tomato, lentils (dry) – 0.02*,**; cabbage (head), Chili pepper (dry) – 0.1*,**; beans (pods, and green seeds) – 3.0*,**; peanuts – 0.5*,**; chicken meat and offal, eggs – 0.03*,**
239	Clethodim	Dry beans – 2.0*,**; cottonseed oil, edible – 0.5*,**; edible offal – 0.2*,**; eggs – 0.05*,**; field peas (dry) – 2.0*,**; sugar beet – 0.1; garlic – 0.5*,**; meat of mammals (except sea mammals) – 0.2*,**; milk – 0.05*,**; onion (bulb) – 0.5; ground nut – 5.0*,**; potato – 0.5; poultry meat and offal – 0.2*,**; rapeseed (seed, oil) – 0.5; soybean beans (dry) – 10.0*; soybean oil for human consumption – 0.5; sunflowerseeds (seeds) – 0.5; sunflowerseed oil (crude) – 0.1; tomatoes – 1.0*,**; carrots, red beet – 0.1 onion, carrot, soya bean, soya bean oil, sugar beet, red beet – 0.1; potatoes, sunflower (seeds and oil) – 0.2; rape (seeds, oil) – 0.5
240	Clefoxydim	Rice -0.05*
241	Clodinafop -propargyl	Cereal grain - 0.05
242	Clozantel	For cattle: fat, kidney-3.0; liver, meat -1.0; for sheep: fat-2.0; meat, liver-1.5; kidney -5.0
243	Cloquintocet-mexyl	Cereal grain-0.1
244	Clomazone	Soya bean, soya-bean oil - 0.01; rice-0.2*; corn (grain) , carrot, sugar beet, rape (seeds, oil) -0.1
245	Clopyralid	Cereal grain-0.2; cabbage -0.05*; corn (grain) -
246	2-ethylhexyl ether of Clopyralid	2.0; meat and meat products -0.3; milk and milk products, wild mushrooms and berries–0.004; corn oil, sugar beet, rape (seeds, oil) - 0.5 NR
247	Clothianidin	Potatoes-0.05; rape (seeds)-0.04; rapeseed oil, sugar beet -0.1; cereal grain – 0.2
248	Clofentezine	Grapes - 2.0; citrus fruits – 0.5**,; seed type fruits – 0.5; potato – 0.05; almond in shell – 5.0*,**; cucumbers – tomato, tree nuts , stone type fruits – 0.5*,**; currants (black, white, red) – 0.2*,**; dry grapes (raisin), strawberry –

		2.0*,**; offal of mammals, eggs, meat of mammals (except sea mammals), milk, poultry meat and offal – 0.05*,**; melons – 0.1*,**; grapes 1.0; Citrus fruits 0.05*,**; fruit (seeded fruit)– 0.5; potatoes 0.05
249	Kresoxim-methyl	Barley – 0.1*,**; cucumbers – 0.5; raisin (dry) – 2.0*,**; mammals' offal, edible – 0.05*,**; grapefruit – 0.5*,**; grapes – 1.0*,**; fat of mammals, except milk fat – 0.05*,**; milk – 0.01*,**; olive oil – 0.7*,**; olives – 0.2*,**; oranges, including hybrids – 0.5*,**; seed type fruits – 1.0 (K); chicken meat – 0.05*,**; cereal grain (wheat, rye) – 0.05*,**; tomatoes – 0.5; berries – 1.0 eucumbers, grapes,; fruit (seeded fruit)–0.2; berries–0.1*
250	Crotoxyphos	Milk, meat products, milk products -0.004; meat - 0.05
251	Coumaphos	Milk products, eggs – 0.01; beef, poultry meat -0.1; pork, meatproducts-0.2
252	Lenacyl	Sugar beet, Red beet-0.1
253	Lindane	Cereal grain (barley, wheat, oats, rye) – 0.01*,**; offal of mammals – 0.01*,**; eggs – 0.01*,**; corn (grain) – 0.01*,**; meat of mammals (except sea mammals) -0.1*,**; milk – 0.01*,**; poultry meat – 0.05*,**; poultry offal – 0.01*,**; sorghum – 0.01*,**; sweet corn – 0.01*,**
254	Luphenuron	Fruits (stone fruits), potatoes - 0.04; tomatoes-0.5; grapes-0.1
255	Lambda-cygalotrine	Fruits (stone fruits) -0.03*; dry hop-1.0*; mustard (seeds, oil) - 0.1*; rape (seeds, oil), soya (beans, oil) -0.1; corn (grain), cabbage, tomatoes, peas, cereal grain, potatoes, carrot-0.01; fruits (seeded fruits)-0.03; sugar beet, onion -0.02; grapes -0.15
256	Malathion	Apples – 0.5; asparagus – 1.0*,**; beans (dry) – 2.0*,**; beans, except fodder and soya – 1.0*,**; blackberry – 10.0*,**; citrus fruits – 7.0*; cottonseed seeds – 20.0*,**; cottonseed oil for human consumption – 13.0*,**; cucumbers -0.2; grapes – 5.0*; corn – 0.05; leaf mustard – 2.0*,**; pepper – 0.1*,**; pepper Chili (dry) – 1.0*,**; sorghum – 3.0*,**; spinach – 3.0*,**; onion (leaf, bulb) – 5.0; berries (strawberry, currant – black, white, red, gooseberry, raspberry) – 1.0; sweet corn,

		table, boiled in cobs – 0.02*,**; tomato – 0.5; tomato juice – 0.01*,**; wheat – 10.0*; wheat bran, not processed – 25.0*,**; sugar beet, red beet, cabbage, fruits (pomaceous and stone type), melon type, tea – 0.5; peas, soybeans (beans) – 0.3; tobacco, dry hop, mushrooms, groats (except wheat) – 1.0; soybean oil – 0.1; peanuts – 1.0*; bread – 0.3*; mustard, oilseed poppy – 0.1*, animal products – 0.01; sunflowerseed (seeds, oil) – 0.02; potato, carrot – 0.05 cereal grain – 3.0; sugar beet, red beet, fruits (seeded fruits, stone fruits), grapes, cabbage, cucumbers, gourds, tomatoes, tea – 0.5; corn (grain), peas, soya (beans) – 0.3; tobacco, dry hop, mushrooms, grits (other than semolina) – 1.0; soya (oil) – 0.1; peanuts – 1.0*; bread – 0.3*; citrus fruits – 0.2*; mustard, oil poppy – 0.1*; livestock products, berries – 0.01; sunflower (seeds, oil) – 0.02; potatoes – 0.05
257	Maleic gidrazit	Garlic – 15.0; onion (bulb, shallot) – 15.0; potato – 50.0; sugar beet, red beet, carrot, tomato, water melon – 8.0; green tobacco – 30.0
258	Mandipropamid	Broccoli – 2.0*,**; cabbage (head) – 3.0*,**; onion (bulb) – 0.1; potato – 0.5; spring onion – 7.0*,**; pumpkin (summer) – 0.2*,**; pepper – 1.0*,**; pepper Chili (dry) – 10.0*,**; leaf vegetables – 25.0*,**; cucumbers – 0.2*,**; tomato – 1.0; cherry – 20.0*,**; grape – 2.0*,**; raisin (all types) – 5.0*,**; melon – 0.5*,** potatoes – 0.5; tomatoes – 1.0; onion – 0.1
259	Mankozeb	Potatoes, onion, tomatoes, grapes, cucumbers – 0.1
260	Industrial (vaseline) oil -8A	All plant products - RNR
261	Petroleum oil (inhibite)	NR
262	Bis copper (8- oxyquinolate)	Cereal grain, potatoes, fruits (seeded fruits), tomatoes - 1.0; sugar beet - 0.1; grapes - 0.5
263	Copper-bearing substances: -copper hydroxide -copper sulfate –copper oxychloride - copper tricaptolactam dichloride monohydrate (copper check)	Potatoes-2.0; dry hop-10.0*; eggs, meat - 2.0; fruits (seeded fruits, stone fruits), tomatoes, berries, grapes, sugar beet, cucumbers, onion, vegetables, gourds – 5.0; citrus fruits – 20.0 citrus fruits – 5.0
264	Copper tricaptolactam dichloride monohydrate (captolactam part of the molecule)	Sugar beet - 0.5; tomatoes, onion, carrot, apples, grapes -0.15; potatoes-1.0
265	Mesosulfuron - methyl	Cereal grain -0.5

266	Mesotrione	Corn (grain, oil)-0.1
267	Mecoprop	Cereal grain - 0.25
268	Menazon	Fruits (seeded fruits, stone fruits), vegetables, gourds, potatoes, sugar beet, legumes, tobacco - 1.0
269	Mewpiquat chloride	Rapeseed (seeds, oil) – 3.0
270	Metazachlor	Cabbage - 0.02; mustard (seeds) -0.02*; mustard (oil), rape (seeds, oil) - 0.1
271	Metazine	Potatoes - 0.05*; peas - 0.1*
272	Metaldehyde	Cereal grain , fruits (stone fruits, seeded fruits), vegetables (other than potato), grapes -0.7; citrus fruits -0.2*; berries- 0.8
273	Metam	NR
274	Metamidofos	Artichoke – 0.2*,**; beans, including fodder and soya beans – 1.0*,**; cottonseed seeds – 0.2*,**; mammals’ offal – 0.01*,**; eggs – 0.01*,**; meat of mammals (except sea) - 0.01*,**; milk – 0.02*,**; potato – 0.05*,**; poultry meat - 0.01*,**; poultry offal – 0.01*,**; soybeans, dry – 0.1*,**; sugar beet - 0.02*,**
275	Metamitron	Sugar beet, red beet - 0.03
276	Metanitrofenilgid razonomezoksalevoy acid-diethyl ether	Cereal grain - 0.1*; cucumber s- NR
277	Metaflumizone	Brussels sprouts – 0.8*,**; chines cabbage – 6.0*,**; mammals’ offal – 0.02*,**; eggplants – 0.6*,**; salad – 7.0*,**; meat of mammals (except sea) – 0.02*,**; milk fat – 0.02*,**; milk – 0.01*,**; pepper – 0.6*,**; pepper Chilian, dry – 6.0*,**; potato – 0.02*,**; tomato – 0.6*,**
278	Methidathion	Almonds – 0.05*,**; apples – 0.5*,**; artichoke – 0.05*,**; dry beans – 0.1*,**; cabbage (head) – 0.1*,**; CATTLE fat – 0.02*,**; cherry – 0.2*,**; cottonseed seed – 1.0*,**; cottonseed oil, refined – 2.0*,**; cucumbers – 0.05*,**; offal of CATTLE, pigs, sheep – 0.02*,**; eggs – 0.02*,**; goat fat – 0.02*,**; goat meat – 0.02*,**; goat offal edible – 0.02*,**; grapefruit – 2.0*,**; grape – 1.0*,**; hop dry – 5.0*,**; lemon – 2.0*,**; corn 0.1*,**; mandarins – 5.0*,**; meat of CATTLE, pigs, sheep – 0.02*,**; milk – 0.001*,**; nectarines – 0.2*,**; olives – 1.0*,**; onion (bulb) - 0.1*,**; oranges – 2.0*,**; pears – 1.0*,**; peas (dry) – 0.1*,**;

		pig fat – 0.02*,**; pineapple – 0.05*,**; plums – 0.2*,**; potato – 0.02*,**; poultry meat – 0.02*,**; poultry fat – 0.02*,**; poultry offal, edible – 0.02*,**; radish 0.05*,**; rapeseed – 0.1*,**; sheep fat – 0.02*,**; sorghum – 0.2*,**; sugar beet – 0.05*,**; sunflowerseed (seeds) – 0.5*,**; tea, green and black (dried and fermented) – 0.5*,**; tomato – 0.1*,**; walnuts – 0.05*,**
279	Methyl bromide (nonorganic bromide check)	Tomato – 3.0; cucumbers – 2.5; salad 2.5*; dill, celery, parsley - 1.5*; eggplant, pepper – 2.0*; bread and other prepared grain products, cocoa products , dry fruits, milled grain products , peanuts, tree nuts (MRLs at sales and for direct consumption); cocoa beans, cereal grains - 5.0*, dry fruits – 2.0*; milled grain products – 1.0*; peanuts, tree nuts – 10.0* (MRLs for imported products and for grain before milling, after 24 hours of ventilation) - cereal grain, cacao beans (for imported ones after 24 hours of aeration) 50.0; tomatoes 3.0; grain mill stock meant for cooking 10.0; cucumbers 2.5; lettuce 2.5*; dill, celery, parsley 1.5*; egg plants, pepper 2.0*; dried fruit, peanuts, nuts, cocoa products (for direct consumption) 0.5; dried fruit (applied to imported ones after 24 h of aeration) 20.0; peanuts, nuts (applied to imported ones after 24 h of aeration) 100.0
280	Methylisothiocyanate	Cucumbers, tomatoes- 0.05
281	Methiocarb	Artichoke – 0.05*,**; barley – 0.05*,**; Brussels sprouts – 0.05*,**; headed cabbage 0.1*,**; cauliflower – 0.1*,**; hazel nut – 0.05*,**; leek – 0.5*,**; headed salad – 0.05*,**; corn – 0.05*,**; melon – 0.2*,**; onion bulb – 0.5*,**; peas (dry) – 0.1*,**; peas/beans (not ripened) – 0.1*,**; sweet pepper, including clove pepper – 2.0*,**; potato – 0.05*,**; rapeseed (seeds) – 0.05*,**; strawberry – 1.0*,**; sugar beet – 0.05*,**; sunflowerseed – 0.05*,**; wheat – 0.05*,**; wheat straw and hay (dry) – 0.05*,**
282	Metconazole	Rape (grain, oil)- 0.15; grain of cereals – 0.2
283	Metobromuron	Potatoes - 0.1; tobacco – 0.5
284	Metoxychlor	Potatoes - 0.3
285	Metoxuron	Cereal grain, vegetables (other than potato) -

		0.1; carrot – 0.02
286	S- metolachlor	Gourds, cucumbers- 0.05*; tobacco, dry hop- 1.0*; cottonseed (oil) soya (oil), cabbage - 0.02; corn (grain), soya (beans), sunflower (seeds), red beet, rape (grain, oil)-0.1; sunflower (oil), sugar beet -0.05; corn (oil) – 0.1
287	Methoxyfenozone	Peanuts – 0.03*, **; peanut butter edible – 0.1*, **; papaya, grapes – 1.0*, **; avocado, citrus fruits, cranberry – 0.7*, **; carrots, beans dry – 0.5*, **; beans shelled – 0.3*, **; corn, sweet corn, cobs – 0.02*, **; beans (pods whole, seeds), dry grapes (all types f raisin) – 2.0*, **; broccoli – 3.0*, **; blueberry – 4.0*, **; peas (dry) – 5.0*, **; apple pure (dry), headed cabbage, cottonseed seeds – 7.0*, **; celery, salad headed – 15.0*, **, leaf salad, leaf mustard – 30.0*, **; offal of mammals, eggs – 0.01*; fat of mammals (except milk fat), meat of mammals (except sea mammals) – 0.2*, **; milk – 0.05*, **
288	Methomyl	Fruits (seeded fruits) (apples, pears), grape – 0.3; beans (dry) – 0.05*, **; citrus fruits – 1.0, pulp of citrus fruits (dry) – 3.0*, **; vegetables with edible fruits, pumpkin type vegetables – 0.1*, **; cottonseed (seed, small, milled, edible) – 0.05*, **; cottonseed (oil, edible) – 0.04*, **; cottonseed (seeds), salad headed and leaf, stone type fruits (peaches, nectarines), soya beans (dry), soya oil – 0.2*, **; beans (dry) – 0.05*, **; beans (except broad beans and soybeans, ordinary beans (pods and seeds) – 1.0*, **; soya beans, onion (bulb), plums – 1.0*, **; soybean flour – 20.0*, **; corn (seeds, oil), potato, oats – 0.02*, **; mint (dry) – 0.5*, **; peas (pods and seeds) – 5.0*, **; oats, pepper – 0.7*, **; Chili pepper (dry) – 10.0*, **; rapeseed (seeds), asparagus, wheat, wheat sprouts, barley – 2.0*, **; wheat bran, not processed – 3.0*, **; wheat flour – 0.03*, **; meat and offal of mammals (except sea mammals), poultry meat and offal, eggs, milk – 0.02*, ** - Fruits (seeded fruits) (apples, pears), grape – 0.2; grapes – 0.05
289	Methoprene	Grain of cereals – 10.0*, **; wheat bran, not processed – 25.0*, **; corn oil (crude) –

		200.0*,**; meat of mammals (except sea) - 0.2*,**; milk - 0.1*,**; poultry meat, eggs and offal, offal of mammals - 0.02*,**
290	Metrafenone	Grain of cereals - 0.5; grapes - 5.0
291	Metribuzin	Tomatoes, potatoes-0.25; soya (beans, oil), corn (grain)- 0.1
292	Metsulfuron- methyl	Cereal grain, millet -0.05
293	Mefenoxam (metalaxyl, metalaxyl M)	Potato, sugar beet, red beet - 0.05; cucumbers, tomato, cabbage (headed) - 0.5; dry hop - 10.0*; sunflowerseed (seeds, oil), corn (grain), rapeseed (seeds, oil), grain of cereals - 0.1; onion (bulb) - 2.0; grapes - 2.0; tobacco - 1.0*; cauliflower, broccoli, gherkins - 0.5*,**; spinach - 2.0*,**; avocado, Brussels sprouts, cocoa beans, pumpkin, melon, water melon, currant (red and black) - 0.2*,**; citrus - 5.0*,**; carrots, cottonseed (seed); peas fresh, shelled, soya beans (dry) - 0.05*,**; salad headed - 2.0*,**; peanuts, pepper, pomaceous fruits - 1.0*,**; Chili pepper (dry) - 10.0*,** potatoes, onion, sugar beet, red beet - 0.05; cucumbers, tomatoes - 0.5; cabbage - 0.01; dry hop - 5.0*; tobacco - 1.0*; sunflower (seeds, oil), grapes, corn (grain), rape (grain, oil); cereal grain - 0.1
294	Mefenpyr-diethyl	Cereal grain, corn (grain, oil) - 0.5
295	Miclobutanil	Banana, dry hop, stone type fruits - 2.0*,**; grapes - 1.0*,**; currant black, pomaceous fruits, prunes - 0.5*,**; tomato - 0.3*,**; plums, including prunes - 0.2*,**; strawberry - 0.1*,**; meat and offal of CATTLE and poultry, eggs, milk - 0.01*,** NR
296	Milneb	Plant food products -1.0
297	Molinat	Rice - 0.2
298	Monolinuron	Potatoes - 0.02; cereal grain, grain legumes - 0.2
299	Naled	Vegetables-0.1; meat -0,3; potatoes, eggs, milk and milk products -0.2
300	Napropamide	Sunflower (seeds) - 0.15*; sunflower (oil) - 0.05*; tomatoes, cucumbers, marrows, pumpkin -0.1*; tobacco -1.0*
301	Sodium silicofluoride	Meat (including natural background) -0.4
302	Sodium salycilate	NR
303	Sodium trichloroacetate	Berries-0.01; sugar beet, red beet, vegetables (other than potato), fruits (seeded fruits, stone fruits), sunflower (seeds, oil), cereal grain,

		grain legumes -0.01
304	Naftalen-1- Ilthiocarbamide	NR
305	Naphthalic anhydride	Cereal grain -0.02
306	Neonol	NR
307	Nicosulfuron	Corn (grain)- 0.2; corn (oil)-0.1
308	Nitroalkilfenolates	NR
309	Nitrotrichloro-methane	Grain to be processed
310	Novaluron	Apple cake, dry – 40.0*,**; cottonseed seeds – 0.5*,**; mammals' offal, edible – 10.0*,**; meat of mammals (except sea) – 10.0*,**; milk fat – 7.0*,**; milk – 0.4*,**; seed type fruits – 3.0*,**; potato – 0.01*,**; poultry meat – 0.01*,**; poultry offal – 0.01*,**; soybean beans, not ripened – 0.01*,**; tomato – 0.02*,**
311	Nonylphenol	NR
312	Nore	Plant food products -0.1
313	Oxadixyl	Potatoes -0.1; wet hop - 0.25; grapes, tomatoes -0.5; sugar beet - 1.0*; fruits (seeded fruits) - 0.5*; tobacco, cucumbers, onion - 0.04
314	Oxamil	Sugar beet – 0.1*; dry hop – 1.0*; tomato, cucumbers – 2.0*; peanuts – 0.05*,**; potato, carrots – 0.1*,**; cottonseed seeds – 0.2*,**; melon, pepper sweet (including clove pepper) – 2.0*,**; citrus fruits – 5.0*,**; meat of mammals (except sea mammals), offal of CATTLE. Goats, horses, pigs and sheep, milk, poultry meat, offal and eggs – 0.02*,** tomatoes, cucumbers – 0.5*; sugar beet – 0.1*; dry hop – 1.0*
315	Oxydemeton-methyl	Barley – 0.02*,**; CATTLE meat – 0.05*,**; all beans, dry – 0.1*,**; cauliflower – 0.01*,**; cottonseed seed – 0.05*,**; eggs – 0.05*,**; cabbage – 0.01*,**; kohlrabi – 0.05*,**; lemon – 0.2*,**; meat of CATTLE, pigs, sheep – 0.05*,**; milk – 0.01*,**; pears – 0.05*,**; pigs fat – 0.05*,**; potato – 0.01*,**; poultry fat – 0.05*,**; poultry meat – 0.05*,**; rye – 0.02*,**; sheep fat – 0.05*,**; sugar beet – 0.01*,**; wheat – 0.02*,**
316	Oxicarboxin	Cereal grain 0.2*
317	Oximethylethyl ketone	NR
318	Oxyfluorfen	Fruits (seeded fruits), onion, sunflower (seeds, oil)- 0.2
319	Oleic alcohol (HD-OCENOL)	NR

320	Paraquat	Tea (green and black) (fermented and dry) – 0.2*,**; leaf type vegetables – 0.07*,**; sorghum – 0.003*,**; dry hop, olives – 0.1*,**; berries and other small type fruits, seed and stone type fruits – 0.0a*,**; citrus fruits, vegetables with edible fruits, pumpkin type – 0.02*,**; sunflowerseeds (seeds), cottonseed seeds – 2.0*,**; legumes – 0.5*,**; corn – 0.03*,**; tree nuts, corn flour, vegetables with edible fruits, except pumpkin type, rice – 0.05*,**; vegetables with edible roots and tubers, poultry and mammals' meat and offal (except sea mammals), eggs, milk – 0.005*,**
321	Parathion-methyl	Fruits (seed type) – 0.2; tomato – 0.002; pea, grain of cereals – 0.1; sugar beet – 0.05; dry peas – 0.3*; stone type fruits (nectarines, peaches) – 0.3*,**; potato, beans (dry), cabbage (headed) – 0.05*,**; grapes – 0.5*,**; dry grapes (all kinds of raisin) – 1.0*,** fruits (seeded fruits) – 0.004; tomatoes – 0.002; peas, cereal grain – 0.1; sugar beet – 0.05
322	Pebulat	Vegetables (other than potato), sugar beet - 0.05; tobacco -0.1
323	Pendimethalin	Soya (beans, oil), garlic, tobacco, dry hop - 0.1*; tomatoes, carrot , cucumbers-0.05*; onion, parsley, cabbage, cottonseed (oil) - 0.05; sunflower (seeds, oil)-0.1; carrot – 0.2
324	Penconazole	Cucumbers, water melon – 0.1; grapes – 0.3; tomato – 0.2*; fruits (seed type0, melon – 0.2; grapes, fruits (stone type), except nectarines and peaches) – 0.3; grain of cereals – 0.005; berries – 0.1; dried grapes (all kinds of raisin), dry hop – 0.5*,**; nectarines, peaches, meat and offal of Cattle, poultry meat and eggs – 0.05*,**; milk – 0.01*,** cucumbers, berries, water melons – 0.1; tomatoes – 0.1*; fruits (seeded fruits), melons – 0.2; grapes, fruits (seeded fruits) – 0.3; cereal grain – 0.005
325	Penoxsulam	Rice -0.5
326	Pentachlor	Tomatoes -1.5
327	Pencycuron	Potatoes-0.1
328	Permethrin	Almonds – 0.1*,**; asparagus – 1.0*,**; beans (dry) – 0.1*,**; hop (dry) – 50.0*,**; horse radish 0.5*,**; cabbage (headed, Savoy, Chinese) – 5.0*; broccoli – 2.0*,**; kohlrabi –

		<p>0.1*,**; leek – 0.5*,**; salad (headed) – 2.0*,**; cucumbers (including gherkins) – 0.5*,**; tomato – 1.0*; potato – 0.05; carrot – 0.1*,**; sugar beet – 0.05; pepper – 1.0*; cauliflower – 0.5*; celery – 2.0*,**; eggplant – 1.0*,**; spinach – 2.0*,**; spring onion – 0.5*,**; radish – 0.1*,**; citrus fruits – 0.5*,**; kiwi – 2.0*,**; gooseberry – 2.0*,**; strawberry, dewberry- 1.0*; grapes – 2.0*; melon – 0.1; pumpkin – 0.5*,**; grain of cereals – 2.0; sunflowerseed (oil for human consumption, crude) – 1.0; sweet corn (grain) – 0.1; soybeans (beans, dry) – 0.05; soybean oil, crude – 0.1; coffee (beans) – 0.05*,**; beans (whole pods and/or not ripened grain) – 1.0**, rapeseed (seed) – 0.05*,**; cottonseed (seeds) – 0.5*,**; cottonseed oil for human consumption – 0.1; meat of mammals (except sea mammals) – 1.0*,**; eggs – 0.1*,**; offal of mammals – 0.1*,**; poultry meat – 0.1*,**; mushrooms – 0.1*,**; olives – 1.0*,**; peanut – 0.1*,**; peas (shelled, fresh) – 0.1*; Chili pepper (dry) – 10.0*,**; pistachios – 0.05*,**; fruits (seed type) – 2.0*; fruits (stone type) – 2.0*; green and black tea (fermented and dried) – 20.0*,**; wheat bran – 5.0*,**; wheat flour – 0.5*,**; wheat sprouts – 2.0*,**; wheat flour, wholegrain – 2.0*,**; rice – 0.01</p> <p>cottonseed (oil), sunflower (oil), soya (oil), corn (grain) – 0.1; fruits (seeded fruits), rice – 0.01; fruits (stone fruits), grapes – 0.01; potatoes – 0.05; melons, cereal grain, cucumbers – 0.1; sugar beet, soya (beans), pea, cabbage – 0.05; sunflower (seeds) – 1.0; pepper, tomatoes – 0.4; berries – 0.2</p>
329	Picoxystrobin	Grain of cereals – 0.2
330	Pinoxaden	Grain of cereals-1.0
331	Pinolene	NR
332	Picloram	Cereal grain, corn (grain), rape (grain, oil) – 0.01; wild berries -0.5
333	Piperonyl butoxide	Grain of cereals – 30.0*,**; citrus – 5.0*,**; juice of citrus fruits – 0.05*,**; dried fruits, legumes – 0.2*,**; vegetables with edible fruits, pumpkin type, peanuts (in shell) – 1.0*,**; pepper, tomato – 2.0*,**; root type vegetables (except carrots) – 0.5*,**; tomato

		juice – 0.3*,**; pepper Chili (dry) – 20.0*,**; leaf salad, leaf mustard, spinach – 50.0*,**; corn (oil), wheat bran – 80.0*,**; kidney of Cattle – 0.3*,**; meat of Cattle – 5.0*,**; poultry meat – 7.0*; liver of Cattle, goats, pigs, sheep, eggs – 1.0*,**; kidney of goats, pigs, sheep (except kidney of Cattle), milk of Cattle. – 0.2*,**; meat of mammals (except sea mammals) – 2.0*,**; milk (except milk of Cattle.) – 0.05*,**; poultry offal – 10.0*,**
334	Pirazosulfuron-ethyl	Rice-0.1
335	Pirazofos	All food products – 0.01
336	Pyraclostrobin	Grapes -2.0; fruits (seed type) – 0.5; grain of cereals – 0.5; corn (grain and oil), soybean oil – 0.02; soybean (beans0 – 0.05; almonds in shell, salad (headed), raspberry (red, black) – 2.0*,**; almond shelled, bananas, peanuts (in shell), peas (pods, not-ripened seeds0, pecan, potato – 0.02*,**; beans 9dry), cabbage (headed), cantaloupe (melon), onion (bulb), sugar beet – 0.2*,**; blueberry, citrus fruits, cabbage (orchard), pistachios, fruits (stone type) – 1.0*,**, brussel sprouts, coffee beans, eggplants, peas (dry), pumpkin (ordinary), sunflowerseed (seeds), tomato – 0.3*; carrots, cucumbers, lentils (dry), meat of mammals (except sea mammals), pepper, radish, strawberry – 0.5*,**; dried grapes (raisin) – 5.0*,**; offal of mammals, poultry meat and offal, eggs, garlic, mango, papaya – 0.05*,**; broccoli, Chinese cabbage and cauliflower – 1.0*,**; hop (dry) – 15.**,**; leek – 0.7*,**; milk – 0.03*,** fruits (seeded fruits)-0.3; cereal grain-0.1
337	Pyrethrins	Grain of cereals – 0.3*,**; legumes – 0.1*,**; citrus fruits, peppers, vegetables with edible roots and tubers, tomato, vegetables with edible fruits, pumpkin type vegetables – 0.05*,**, dry fruits – 0.2*,**; peanuts, pepper Chili (dry), tree nuts – 0.5*,**
338	Pyridaben	Fruits (seeded fruits) – 0.2; citrus fruits - 0.3
339	Pyridat	Corn (grain)-0.05
340	Pyridafention	Cabbage -0.1; sugar beet, citrus fruits (pulp) - 0.1 *
341	Pyrimethanil	Almonds, onion (bulb) – 0.2*,**; apple puree (dry) – 40.0*,**; apricot, strawberry, beans

		(pods and/or not-ripened seeds), salad (headed type), Welsh onion – 3.0*,**; nectarines, cherry, grapes – 4.0*,**; fruits (seed type), citrus fruits – 7.0*,**; plums – 2.0*,**; bananas – 0.1*,**; carrots – 1.0*,**; tomato – 0.7*,**; nuts – 0.5*,**; dry grapes (all types of raisin) – 5.0*,**; potato, meat of mammals (except sea mammals) – 0.05*,**; milk, offal of mammals – 0.1*,**
342	Pyrimicarb	Fruits (seeded fruits, stone fruits) – 0.05; cucumbers-0.1; dry hop- 1.0*; potatoes, sugar beet, cottonseed (oil), pea - 0.02; fruits (seed type) – 2.0**, fruits (stone type) – 5.0**, berries, except strawberry – 1.0**, strawberry – 3.0**, asparagus – 0.01*,**; vegetables with edible roots and tubers, grain of cereals, rapeseed (seeds), sweet corn (boiled in cobs) – 0.05*,**; garlic, onion (bulb), sunflowerseed (seeds) – 0.1*,**; melon, corn (grain), beans, legumes (dry), except soybeans – 0.2*,**, cabbage – 0.3*,**; vegetables with edible fruits, except pumpkin type – 0.5*,**, bean type vegetables, except soybeans – 0.7; grapes and other small size fruits, vegetables with edible fruits, pumpkin type vegetables, except melon and water melon – 1.0*,**, citrus fruits – 3.0*,**, salad (headed type) and leaf type, artichoke – 5.0*,**, Chili pepper (dry) – 20.0*,**, meat of mammals (except sea mammals); poultry meat, offal and eggs, milk – 0.01*,**
343	Pirimiphos-methyl	Berries, cultured mushrooms - eggs – 0.004; melons, peppers, egg-plants, sugar beet -0.2*; Russian turnip, turnip, cabbage, celery (green), fruits (stone fruits), grapes, tea -0.5*; citrus fruit (pulp) - 0.1*; potatoes, radish, celery (celiac), carrot -0.05*; rice, tobacco - 1.0*; tomatoes, cucumbers- 0.2; eggs – 0.01; grain of cereals – 7.0; wheat bran, not processed – 15.0*,** poultry meat - 0.1; poultry liver -0.5; meat of mammals (except sea mammals), offal of mammals, poultry offal, except liver, milk – 0.01*,** pea 5.0*; cereal grain – 0.1;
344	Pirimiphos-ethyl	Corn (grain) -0.1
345	Pyriproxyfen	Fruits (seeded fruits), cucumbers, tomatoes - 0.2; citrus fruits – 0.5*,**, cottonseed (seeds) –

		0.05*,**; cottonseed (oil) – 0.01*,**; meat and offal of Cattle and goats – 0.01*,**
346	Pyroxsulam	Grain of cereals – 0.5
347	Poly-beta- hydroxybutyric acid	RNR
348	Polyhexamethylene guanidine	Potatoes - 0.2
349	Polyoxyethylene dodecyl ether	NR
350	Pirimisulfuron	Corn (grain)-0.05
351	Products of metabolism of ginseng endophyte fungi	RNR
352	Products of metabolism of sea-buckthorn endophyte fungi	RNR
353	Progeksadion calcium	Fruits (seed type) – 0.5
354	Proquinazid	Grapes-0.5
355	Prometryn	Caraway seeds -0.1*; sunflower (seeds, oil), coriander, soya (beans, oil), pea, garlic, kidney beans, potatoes, lens, corn (grain, oil) -0.1; carrot, potato, celery, fennel, parsley -0.02
356	Propazine	Sorghum, coriander - 0.2*; cereal grain, grain legumes -0.2; carrot – 0.04
357	Propaquizapop	Cottonseed (oil), flax - 0.01; sugar beet , rape (grain, oil)-0.1; cabbage -0.2
358	Propamocarb hydrochloride	Cucumbers, potatoes-0.1; salad (headed and leaf) -15.0**, radish -1.0**, potato – 0.3; tomato, cucumbers – 10.0; cauliflower - 0.2*,**; eggplants – 0.3*,**; spinach – 40.0*,**; pepper Chili (dry) – 10.0*,**; pepper sweet, including clove pepper – 3.0*,**; chicory (sprouts) – 2.0*,**; meat and offal of mammals (except sea mammals) and poultry, milk, eggs – 0.01*,**
359	Propanil	Rice-0.3
360	Propargite	Soya (beans, oil)-0.1; cottonseed (oil) -0.1*, cucumbers-0.2*; fruits (stone fruits) -0.5* - 4.0; fruits (seeded fruits) – 3.0*; apple juice – 0.2*; citrus fruits – 3.0*, 0.3*; citrus fruits pulp, dry – 10.0*,**; almonds – 0.1*,**; beans (dry) – 0.3*; cottonseed (seeds) – 0.1*,**; grapes – 7.0* 0.2; chicken peas, dry – 0.3*; cottonseed (seeds) – 0.1*,**; grapes – 7.0*; grape juice – 1.0*,**; grapes dry (all types of raisin) – 12.0*,**; offal of mammals – 0.1*,**; eggs – 0.1*,**; dry hop – 100.0* 30.0; corn – 0.1*,**; corn flour – 0.2*,**; corn oil (crude) – 0.7*,**; corn oil (for human consumption – 0.5*,**; peanuts, milk, meat and offal of mammals (except sea mammals) and of

		poultry, eggs – 0.1*,**; peanut butter for human consumption – 0.3*,**; potato – 0.03*,**; tea green and black (black tea fermented and dried) – 5.0*,**; tomato – 2.0*,**
361	Propachlor	Cabbage, onion, garlic, Russian turnip, turnip - 0.2; cereal grain, grain legumes -0.3; corn - 0.3*; soya (beans) – 0.1
362	Propizamid	Sugar beet - 0.1; endive - 1.0*
363	Propisochlor	Corn, rapeseed (seeds, oil), sunflowerseed (seeds, oil) – 0.1
364	Propetamphos	Meat-0.02; milk -0.01
365	Propiconazole	Cereal grain (except barley), sugar beet, rape (grain, oil)- 0.1; barley – 0.2, red beet, berries (except cranberry)- 0.05, cranberry – 0.3; grapes-0.5; banana – 0.1*,**; coffee (beans), pecan, pineapple, sugar cane – 0.02*,**; meat and offal of mammals (except sea mammals), poultry meat, eggs, milk – 0.01*,**; corn, popcorn, sweet corn (table, boiled in cobs) – 0.05*,**; soybean (beans) – 0.07*,**
366	Propocsure	Livestock products – 0.01
367	Prosulfocarb	Potato – 0.1
368	Prosulfuron	Corn (grain)-0.02; cereal grain , millet - 0.05
369	Protioconazole (after protioconazole destio) protioconazole destio (basic metabolite of active ingredient of procioconazole)	Cereal grain (barley, wheat, rye – 0.3; oats – 0.5*; rapeseed (seed) – 0.1; rapeseed (oil) – 0.05; sugar beet – 0.3*,**; peanut – 0.02*,**; prunes – 1.0*,**; meat of mammals (except sea mammals) – 0.01*,**; offal of mammals – 0.5*,**; rapeseed (seed, oil) – 0.05; cereal grain – 0.3
370	Prothiofos	Cottonseed (oil), grapes -0.1; cabbage - 0.05*
371	Profenfos	Cottonseed seeds – 3.0*,**; offal of mammals – 0.05*,**; eggs – 0.02*,**; mango – 0.2*,**; meat of mammals (except sea mammals) – 0.05*,**; milk – 0.01*,**; pepper Chilean – 5.0*,**; pepper Chilean (dry) – 50.0*,**; poultry meat and offal – 0.05*,**; tea (including herbal tea) – 0.5*,**; tomato – 10.0*,**; cabbage, onion, garlic, Russian turnip, turnip - 0.2; cereal grain, grain legumes -0.3; corn -0.3*; soya (beans) – 0.1
372	Prochloraz	Sugar beet – 0.1; cereal grain – 2.0; citrus fruits – 10.0*,**; flax seeds – 0.05*,**; mushrooms – 3.0*,**; pepper (white and black) – 10.0*,**; sunflowerseeds (seeds) – 0.5*,**;

		sunflowerseed (oil) – 1*,**; rapeseed (seeds) – 0.7*,**; bran, not processed – 7.0*,**; offal of mammals – 10.0*,**; meat of mammals (except sea mammals) – 0.5*,**; milk – 0.05*,**; poultry meat – 0.05*,**; poultry offal – 0.2*,**; eggs – 0.1*,** cereal grain – 0.05; sugar beet – 0.1
373	Procymidone	Cucumbers – 2.0*; tomato, grapes – 5.0*; peas – 1.0*; peas (new pods) – 3.0*,**; legumes (whole pods or/and not ripened seeds) – 1.0*,**; cabbage (headed), plums, peach, gherkin – 2.0*,**; raspberry, strawberry, cherry – 10.0*,**; pear – 1.0*,**; sunflowerseed (seeds), onion (bulb) – 0.2*,**; sunflowerseed (oil) – 0.5*,**; salad (headed), pepper – 5.0*,**; pepper Chili (dry) – 50.0*,** cucumbers, tomatoes, grapes – 0.5*; peas – 1.0*
374	Rimsulfuron	Corn (grain), potatoes -0.01; corn (oil)-0.02; tomato – 0.05
375	Sulfur	RNR
376	Carbon sulphide(product of sulfur block combustion)	RNR
377	Sethoxydim	Sugar beet, soya (beans, oil) - 0.1; citrus fruits, carrot -0.02; fruits (seeded fruits, stone fruits), grapes- 0.05*; cabbage - 0.03
378	Simazine	Cereal grain, corn (grain), potatoes, cabbage - 0.1; fruits (seeded fruits, stone fruits)-0.2; citrus fruits -0.05*; tea, grapes - 0.01; berries (including wild berries) –0.02
379	Mixture of non-ionic surfactants of fixed composition (Amigo adjuvant, KS)	NR
380	Mixture of non-ionic surfactants of fixed composition (PAVDASH)	NR
381	Mixture of non-ionic surfactants in Corvette	NR
382	Spinetoram	Salad (headed and leaf) – 10.0*,**; oranges (including hybrids) – 0.07*,**; fruits (seed type) – 0.05*,**; tomato – 0.06*,**; sugar beet, tree nuts – 0.01*,**; meat of mammals (except sea) – 0.2*,**; offals of mammals, milk – 0.01*,**; milk fat – 0.1*,**
383	Spinosad (Spinosin A + Spinosin D)	Cucumbers – 1.0; pepper – 2.0; potato – 0.5; almond in shell – 0.01*,**; apples – 0.1*,**; celery – 2.0*,**; grain cereal – 1.0*,**; citrus fruits – 0.3*,**; cottonseed seed – 0.01*,**;

		cottonseed oil, for food consumption – 0.01*,**; grape – 0.5*,**; dry grape (all types of raisin) – 1.0*,**; kiwi – 0.05*,**; leaf vegetables – 10.0*,**; soybean (beans, dry) – 0.01*,**; pepper Chili (dry) – 3.0*,**; fruits (stone type) – 0.2*,**; tomatoes – 0.3*,**; wheat bran, not processed – 2.0*,**; cabbage (head type, kale buds) – 2.0*,**; kidney of Cattle – 1.0*,**; liver of Cattle – 2.0*,**; meat of Cattle – 3.0*,**; milk of Cattle – 1.0*,**; meat of mammals (except sea) – 2.0*,**; milk fat of Cattle – 5.0*,**; offal of mammals – 0.5*,**; eggs – 0.01*,**; poultry meat – 0.2*,** cucumbers 0.5* ; pepper 1.0* ; potatoes 0.05*
384	Spirodiclofen	Citrus fruits – 0.4*,**; cucumbers, gherkin – 0.07*,**; currant (red, black, white) – 1.0*,**; dried grape (all types of raisin) – 0.3*,**; papaya, coffee beans – 0.03*,**; pepper, sweet (including Spanish pepper and small peppers), grape – 0.2*,**; seed type fruits – 0.8*,**; fruits (stone types), strawberry – 2.0*,**; tomato – 0.5*,**; hop (dry) – 40.0*; tree nuts, offal of mammals – 0.05*,**; meat of mammals (except sea) – 0.01*,**; milk – 0.004*,**
385	Spiroxamine	Cereal grain - 0.2; grapes-2.0; rice-0.2*; sugar beet -0.1
386	Spirotetramat	Almond in shell – 10.0*,**; hop dry – 15.0**; leaf vegetables – 7.0*,**; cabbage (head type, buds, broccoli, Chinese, cauliflower) – 2.0*,**; celery – 4.0*,**; potato – 0.8*,**; citrus fruits – 1.0**; grapes (all types of raisin) – 4.0*,**; prunes – 5.0*,**; fruits (seed type) – 1.0**; fruits (stone type) – 3.0**; tomato – 2.0**; cucumbers – 0.2**; tree nuts – 0.5*,**; Chili pepper (dry) – 15.0**; pepper (chili and other varieties) – 2.0**; offal of mammals – 0.03*,**; meat of mammals (except sea) – 0.01*,**; milk – 0.005*,**
387	Suplrofos	NR
388	Monoethanolamine salt of sulfanilic acid	Cereal grain -1.0
389	Sulfometuron-methyl	NR
390	Sulfometuron- methyl potassium salt	NR
391	Sulphuryl fluoride	Grain of cereals – 0.05*,**; bran of grain crops, processed and not-processed (except

		buckwheat), wheat flour, rye flour, rye flour whole grain. Whole grain wheat flour, corn flour, corn groats, rice polished, rice milled, wheat sprouts - 0.1*,**, dried fruits - 0.06*,**, tree nuts - 3.0*,**
392	Tau-fluvalinate	Fruits (seeded fruits), cucumbers, grapes - 0.2; cereal grain, soya (beans, oil) -0.01; fruits (stone fruits) - 0.01*;rape (grain, oil), potatoes-0.1; tomatoes -0.1
393	Tebuconazole	Cereal grain (barley, oat, wheat, rye, etc.), millet, sunflower (seeds, oil)-0.2; grapes - 2.0-1.0;millet - 0.2; sugar beet-0.1; corn (grain), soya (beans. oil)-0.1; rapeseed (seed) - 0.5 rapeseed (seed, oil)-0.3; rice- 2.0; pumpkin - 0.02*,**, tomato - 0.2*,**, bananas - 0.05*,**, cherry - 5.0*,**, coffee (beans) - 0.1*,**, coffee (beans roasted) - 0.5*,**, cucumbers - 0.2*,**, raisin - 3.0*,**, dry hop - 30.0*,**, peach - 1.0*,**, ground nut - 0.05*,**, pepper Chili (dry) - 5.0*,**, pepper sweet (including clove pepper) - 0.5*,**, fruit (seed type) - 0.5*,**, offal of Cattle - 0.05*,**, meat of mammals (except sea) - 0.05*,**, milk - 0.01*,**, poultry meat - 0.05*,**, poutry offal - 0.05*,**, eggs - 0.05*,**
394	Tebufenotsid	Almond - 0.05*,**, blackberry - 3.0*,**, cabbage broccoli - 0.5*,**, cabbage (headed) - 5.0*,**, citrus fruits - 2.0*,**, cranberry - 0.5*,**, raisin - 2.0*,**, offal of mammals - 0.02*,**, eggs - 0.02*,**, grape - 2.0*,**, kiwi - 0.5*,**, leaf vegetables - 10.0*,**, meat of mammals (except sea) - 0.05*,**, milk - 0.01*,**, mint - 20.0*,**, nectarine - 0.5*,**, peach - 0.5*,**, pecan - 0.01*,**, pepper - 1.0*,**, Chili pepper (dry) - 10.0*,**, seed type fruits - 1.0*,**, poultry meat - 0.02*,**, rapeseed (seeds) - 2.0*,**, raspberry - 2.0*,**8*; rice, milled - 0.1*,**, sugar cane - 1.0*,**, tomato - 1.0*,**, walnut - 0.05*,**
395	Tecnazene	Potato - 20.0*,**
396	Temefos	Vegetables (other than potatoes), sugar beet, cottonseed (oil) -0.3; citrus fruits, milk - 0.01*; meat, eggs-1.0
397	Tepraloxydim	Sugar beet -0.5; soya (beans) -5.0; soya (oil) -

		0.2
398	Terbacil	Citrus fruits, fruits (seeded fruits, stone fruits) - 0.05
399	Terbumeton	Fruits (seeded fruits), grapes -0.1; citrus fruits (pulp) - 0.1*
400	Terbutilazin	Fruits (seeded fruits), grapes, citrus fruit (pulp), sunflower (seeds)-0.1; potatoes, sunflower (oil) -0.05; corn (grain, oil) – 0.1
401	Terbutiuron	Mushrooms- 0.1; berries –NR
402	Terbutrin	Cereal grain - 0.1; potatoes -0.1
403	Terbufos	Banana – 0.05 ^{***} ; coffee beans - 0.05 ^{***} ; mammals offal - 0.05 ^{***} ; egg -0.01 ^{***} ; corn - 0.01 ^{***} ; mammals meat (except marine mammals) - 0.05 ^{***} ; milk – 0.01 ^{***} ; poultry - 0.05 ^{***} ; sorghum 0.01 ^{***} ; sugar beet-0.02 [*] ; corn (sugar, boiled in cob) - 0.01 ^{***} tobacco, potatoes - 0.05; sugar beet-0.01[*]; tobacco; potatoes, corn (grain)–0.05
404	Natural terpenoids (blend)	RNR
405	Tetradifon	Vegetables (other than potatoes), gourds, fruits (seeded fruits)-0.7; cottonseed (oil), grapes - 0.1; citrus fruits (pulp) -0.2*
406	Tetrakonazol	Cereal grain - 0.2; sugar beet – 0.05
407	Tetramethyl methylenediamine oxalate	NR
408	Tetrametrine	Meat, by-products, fats, milk-0.2
409	Tetrafluoron	Cottonseed (oil) - NR; cottonseed (seeds) -0.1
410	Tetrachlorinfos	Cabbage, fruits (seeded fruits, stone fruits)-0.8; grapes, berries - 0.01; cottonseed (oil) - 0.1; dry hop -5.0
411	Teflubenzuron	Brussels sprouts - 0.05 ^{***} ; cabbage - 0.02 ^{***} ; plumps – 0.1 ^{***} ; pomaceous fruits - 1.0 ^{***} ; potatoes - 0.05 ^{***}
412	Tefluthrin	Sugar beet, sunflower (seeds, oil), corn (grain, oil)-0.05; potatoes-0.01
413	Tiabendazole	Cereal grain - 0.02; corn (seeds) – 0.2; millet, rice, pea, sunflower (seeds, oil)- 0.2; tomatoes- 0.1*; potatoes- 15.0; citrus – 5.0 ^{**} ; avocado – 15.0 ^{***} ; bananas – 5.0 ^{***} ; mango - 5.0 ^{***} ; mushrooms - 60.0 ^{***} ; papaya - 10.0 ^{***} ; fruit (seeded) - 3.0 ^{***} ; chicory - 0.05 ^{***} ; bovine animals kidneys - 1.0 ^{***} ; bovine animals liver - 0.3 ^{***} ; bovine animals meat - 0.1 ^{***} ; milk of bovine animals - 0.2 ^{***} ; poultry meat - 0.05 ^{***} ; eggs - 0.1 ^{***}
414	Thiacloprid	Fruits (seeded fruits) – 0.7, rape (oil) -0.3; rape (seed) – 0.5; grapes, potatoes-0.02; berries

		and other small fruits -1.0**, almond in shell - 10.0***; cottonseed (seeds), eggs, poultry meat and poultry by-products, rice, tree nuts - 0.02***; cucumbers, pumpkin - 0.3***; mammals by-products, mustard (seeds), seed fruits, tomatoes - 0.5***; kiwi, melons, water melons, winter squash – 0.2***; mammals meat (except sea mammals), wheat – 0.1***; milk - 0.05***; sweet pepper (including bayberry) - 1.0***; Fruits (seeded fruits), rape (grain, oil) - 0.3; grapes - 0.02; berries - 1.0**
415	Thiametoxam	Cereal grain, potatoes, mustard, rape (grain, oil), sugar beet, cucumbers, peas, sunflower (seeds, oil), cabbage, onion -0.05; tomatoes, egg-plants, pepper-0.2; fruits (seeded fruits), currant, grapes -0.1; corn (seed, oil)- 0.05
416	Thiencarbzon-methyl	Corn (grain, oil) – 0.5
417	Thiodicarb	Cottonseed (oil) - 0.5
418	Thiophanate-methyl	Sugar beet, cereal grain - 1.0; persimmon, feijoa -0.2*; cucumbers, fruits (seeded fruits, stone fruits), grapes – 0.5; currant – 0.01
419	Thiociclam	Sugar beet -0.02; potatoes-NR
420	Thiram	Cereal grain – 0.01; potatoes-0.005 all food products -0.01*; corn (grain,oil) – 0.1
421	Thifensulfuron -methyl	Cereal grain, flax (oil) -0.5; corn (grain), soya (beans, oil) -0.02
422	Tolklofos-methyl	Lettuce (cabbage head, leaves) - 2.0***; potatoes – 0.2***; Radish – 0.1***;
423	Topramezon	Corn (grain, oil) – 0.1
424	Tolyfluanid	Fruits (seeded fruits) - 5.0, cucumbers – 1.0, tomatoes- 1.0* berries - 1.0; grapes- 3.0; berry (raspberry, strawberry, blackberry) – 5.0, currant (black, red, white) – 0.5*; tomatoes – 3.0, dry hop - 50.0***; leek- 2.0***; lettuce (cabbage head) - 15.0***; chili pepper (dry) - 20.0***; sweet pepper, including baypepper - 2.0***; fruits (seeded fruits), cucumbers, tomatoes - 1.0* berries - 1.0; grapes - 0.1*
425	Tralkoxydim	Cereal grain - 0.02
426	Triadimenol	Apples- 0.3; cucumbers, tomatoes – 0.1; cereal grain-0.2; grapes – 2.0, sugar beet-0.1; millet – 0.02*; rice – 0.2; pineapple - 3.0***; artichoke - 0.7***; bananas - 1.0***; coffee (beans) - 0.5***; berries: current (red, black, white), strawberry and others - 0.07***; raisins -

		10.0 ^{*,**} ; vegetables fit for human consumption (other than pumpkin) - 1.0 ^{*,**} ; pumpkin - 0.2 ^{*,**} ; chili pepper (dry) - 5.0 ^{*,**} ; mammals by-products (other than sea mammals) - 0.07 ^{*,**} ; mammals meat (other than sea mammals) - 0.02 ^{*,**} ; milk - 0.01 ^{*,**} ; meat, poultry by-products - 0.01 ^{*,**} ; eggs - 0.01 ^{*,**} ;
427	Triadimefon	Apples (seeded) – 0.3*; artichoke - 0.7 ^{*,**} ; bananas - 1.0 ^{*,**} ; cereal grain – 0.5, coffee (beans) - 0.5 ^{*,**} ; current (red, black, white), strawberry and other berries – 0.7*; grapes – 0.1; dry grapes (raisins)- 10.0 ^{*,**} ; mammals by-products - 0.01 ^{*,**} ; eggs - 0.01 ^{*,**} ; fruit-bearing vegetables, other than pumpkin - 1.0 ^{*,**} ; pumpkin - 0.2 ^{*,**} ; melon – 0.05, mammals meat (other than sea mammals) – 0.02 ^{*,**} ; milk - 0.01 ^{*,**} ; chili pepper (dry) - 5.0 ^{*,**} ; pine apple - 3.0 ^{*,**} ; meat, poultry by-products - 0.01 ^{*,**} ; sugar beet - 0.5 ^{*,**} ; tomatoes – 0.5; cucumbers – 0.5; fruits (seeded fruits, stone fruits)- 0.02; , feijoa – 0.02, rice – 0.2 cereal grain, sugar beet, cucumbers, tomatoes – 0.5; melons, fruits (seeded fruits, stone fruits) – 0.05; grapes – 0.1; berries, feijoa – 0.02
428	Triazofos	Cereal grain - 0.05 ^{*,**} ; cottonseed (seed) - 0.2 ^{*,**} ; cottonseed oil crude - 1.0 ^{*,**}
429	Triallat	Grain legumes -0.05*; cereal grain - 0.05
430	Triasulfuron	Cereal grain - 0.1
431	Tribenuron-methyl	Sunflower (seeds, oil)-0.02; cereal grain -0.01
432	Trimorfamid	Cereal grain, cucumbers, fruits (seeded fruits) - 0.2*; grapes -0.1*
433	Trinexopac-ethyl	Cereal grain -0.2
434	Tris (2-ethylhexyl) phosphate (adjuvant)	RNR
435	Triticonazole	Millet, corn (grain)- 0.1; cereal grain -0.04
436	Tritosulfuron	Cereal grain – 0.01
437	Trifenacin (by definition)	RNR
438	Trifloxystrobin	Grapes - 3.0 ^{*,**} ; bananas - 0.05 ^{*,**} ; cabbage (head, Chinese, broccoli, cauliflower) - 0.5 ^{*,**} ; Brussels cabbage - 0.1 ^{*,**} ; carrots - 0.1 ^{*,**} ; citrus - 0.5 ^{*,**} ; sweet pepper - 0.3 ^{*,**} ; tomatoes - 0.7 ^{*,**} ; strawberry - 0.2 ^{*,**} ; leek - 0.7 ^{*,**} ; almonds - 3.0 ^{*,**} ; celery - 1.0 ^{*,**} ; citrus pulp, dry - 1.0 ^{*,**} ; raisins - 5.0 ^{*,**} ; eggs - 0.04 ^{*,**} ; dry hop – 40.0*; bovine animals, goat, swine, sheep kidneys - 0.05 ^{*,**} ; corn - 0.02 ^{*,**} ;

		mammals meat (other than sea mammals) - 0.05 ^{*,**} ; milk – 0.02*, sweet pepper, including baypepper - 0.3 ^{*,**} ; potatoes - 0.02 ^{*,**} ; poultry meat - 0.04 ^{*,**} ; poultry by-products - 0.04 ^{*,**} ; poultry by-products edible - 0.04 ^{*,**} ; rice - 5.0 ^{*,**} ; sugar beet - 0.05 ^{*,**} ; stone fruits - 3.0 ^{*,**} ; melassa - 0.1 ^{*,**} ; tree nuts - 0.02 ^{*,**} ; wheat - 0.2 ^{*,**} ; fruits (seeded fruits)- 0.1;
439	Triflumizol	Cereal grain - 0.05*; cucumbers, tomatoes, fruits (seeded fruits)- 0.1*
440	Triflusulfuron -methyl	Sugar beet - 0.02
441	Trifluralin	Cottonseed (seeds and oil), carrots bunching ripeness, water melon -0.25*; parsley bunching ripeness -0.01; sunflower (seeds), cabbage, tomatoes, cucumbers, garlic, egg-plants, pepper, onion, soya (beans, oil), sunflower (oil), - 0.1; carrot - 0.01 *; tobacco - 0.5; rape (grain, oil)-0.1
442	Triforin	Fruits (seeded fruits) – 2.0; , grapes -0.01*; cucumbers -0.1 ; berries(blueberry, strawberry black currant, gooseberry) - 1.0 ^{*,**} ; stone fruits: cherries, plumps - 2.0 ^{*,**} ; peach - 5.0 ^{*,**} ; tomatoes - 0.5 ^{*,**} ; legumes (pods or uNRipe seeds) - 1.0 ^{*,**} ; vegetable yields fit for consumption, pumpkin family - 0.5 ^{*,**} ; Fruits (seeded fruits), grapes-0.01*; cucumbers-0.1
443	Trichlorfon	Cereal grain, corn (grain), gourds, grapes, leafy vegetables, cabbage, cucumbers, pepper, tomatoes, soya (beans, oil), sunflower (seeds, oil), potatoes, grain legumes, mustard, rice, fruits (seeded fruits, stone fruits)-0.1; sugar beet, onion, carrot, egg-plants, marrows - 0.05; cottonseed (oil) - 0.1*; mushrooms - 0.2; wild berries, milk, milk products, meat products- 0.01
444	Famoxadone	Barley, cucumbers, pumpkin, wheat bran not processed - 0.2 ^{*,**} ; dry grapes (raisins) - 5.0 ^{*,**} ; meat and mammals by-products 9 other than sea mammals) - 0.5 ^{*,**} ; eggs, poultry meat and by-products - 0.01 ^{*,**} ; grapes – 2.0; tomatoes – 1.0; milk - 0.03 ^{*,**} ; potatoes-0.05; wheat - 0.01 ^{*,**} ; onion – 1.0; sunflower (seeds, oil)- 0.1; tomatoes-0.2; grapes-0.25
445	Fenazaquin	Fruits (seeded fruits)-0.2; grapes - 0.01
446	Fenamidone	Potatoes - 0.03; tomatoes - 0.5

447	Fenamiphos	Apples, bananas, head cabbage and Brussels cabbage, melon, cottonseed (seed), peanuts, cottonseed and peanut non refined oil - 0.05 ^{*,**} ; poultry and mammals meat and by-products (other than sea mammals), eggs - 0.01 ^{*,**} ; milk - 0.005 ^{*,**} ;
448	Fenbukonazol	Apricots, peaches - 0.5 ^{*,**} ; bananas, fat, kidneys, liver, bovine animals meat, rape (grain), sunflower (seed), pumpkin - 0.05 ^{*,**} ; barley, cucumber, melon - 0.2 ^{*,**} ; cherries, grapes - 1.0 ^{*,**} ; eggs, milk, poultry meat and by-products, tree nuts - 0.01 ^{*,**} ; fruits seeded, rye, wheat - 0.1 ^{*,**} ;
449	Fenbutatin-oxide	Almond, pecan, walnut, cucumbers - 0.5 ^{*,**} ; bananas, cherries, prunes, strawberries - 10.0 ^{*,**} ; poultry meat and by-products, eggs, mammals meat (other than sea mammals), milk - 0.05 ^{*,**} ; citrus, grapes, seeded fruits - 5.0 ^{*,**} ; citrus pulp (dry) - 25.0 ^{*,**} ; mammals by-products - 0.2 ^{*,**} ; grape dry meal - 100.0 ^{*,**} ; peaches - 7.0 ^{*,**} ; plumps - 3.0 ^{*,**} ; raisins - 20.0 ^{*,**} ; tomatoes - 1.0 ^{*,**} ;
450	Fenarimol	Seeded fruits, grapes - 0.3; apple meal, hops, chili pepper (dry) - 0.5 ^{*,**} ; artichoke for sowing - 0.1 ^{*,**} ; bananas, dry grapes (raisins) - 0.2 ^{*,**} ; bovine meat kidneys, pecan - 0.02 ^{*,**} ; bovine liver, melon - 0.05 ^{*,**} ; cherries, strawberries - 1.0 ^{*,**} ; peaches, sweet pepper (including bay pepper) - 0.5 ^{*,**} ;
451	Fenbutatinoxide	NR
452	Fenvalerate	Cottonseed (oil) refined and non-refined, corn (grain), soya (beans, oil), pea - 0.1*; fruits (seeded fruits), cereal grains - 2.0*, head cabbage - 3.0*; grapes, potatoes - 0.01*; dry hop - 5.0*; fish - 0.0015; currant - 0.03*; beans shelled, milk - 0.1 ^{*,**} ; beans 9 other than feed beans and soya beans), Chinese cabbage, mammals meat (other than sea mammals), tomatoes, berries (other than currant) and other small fruits - 1.0 ^{*,**} ; broccoli, Brussels cabbage and cauliflower, celery, cherry, citrus, head salad, wheat whole flour - 2.0 ^{*,**} ; cottonseed (seed), cucumbers, melon, tree nuts, wheat flour - 0.2 ^{*,**} ; kiwi, peach, chili pepper (dry), non-processed wheat bran - 5.0 ^{*,**} ; peanut in shell, sunflower (seed), sweet corn (boiled in

		cob) - 0.1 ^{***} ; sweet pepper (including bay pepper), pumpkin and large-fruited winter pumpkin, water melon - 0.5 ^{***} ; vegetables with edible roots and bulbs (other than potatoes and celery) - 0.05 ^{***} ; Cottonseed (oil), corn (grain), soya (beans, oil), pea - 0.1* ; fruits (seeded fruits), cabbage - 0.01; grapes, potatoes - 0.01* ; dry hop - 5.0* ; cereal grain - 0.02; fish - 0.0015; currant - 0.03*
453	Phengexamide	Egg plants - 2.0 ^{***} ; pepper - 2.0 ^{***} ; tomatoes - 2.0 ^{**} ; almonds - 0.02 ^{***} ; fruits stoned (apricots, nectarines, peaches) - 10.0 ^{***} ; cherries - 7.0 ^{***} ; plumps (including prunes) 1.0 ^{***} ; berries and other small fruits (strawberries, blackberry, bilberry, black, red, white currant, gooseberry, black and red raspberry, blueberry and other) - 15.0 ^{**} ; cucumbers (including pickling) - 1.0 ^{**} ; pumpkin - 1.0 ^{***} ; raisins - 25.0 ^{***} ; by products and mammals meat (other than sea mammals) - 0.05 ^{***} ; salad (headed and leaf) - 30.0 ^{***} ; milk - 0.01 ^{***}
454	Fenpiroximat	Soya (beans, oil) grapes, seeded fruit (including apples) - 0.3; bovine kidneys, liver - 0.01 ^{***} ; bovine meat - 0.02 ^{***} ; bovine milk - 0.005 ^{***} ; hop (dry) - 10.0 ^{***} ; oranges (including hybrids) - 0.2 ^{***} ;
455	Fenitrothion	Apples - 0.5*; Cereal grain - 6.0*; mammals by-products - 0.05 ^{***} ; eggs - 0.05 ^{***} ; mammals meat (other than sea mammals) - 0.05 ^{***} ; milk - 0.05 ^{***} ; poultry meat - 0.05 ^{***} ; rice - 0.3; soya beans dry - 0.01 ^{***} ; bread, sunflower (seeds, oil), fruits (stone fruits, pulp), tobacco, sugar beet - 0.1; tea - 0.5*; wild berries and mushrooms - 0.01* Cereal grain - 1.0; rice - 0.3; bread, sunflower (seeds, oil), fruits (seeded fruits, stone fruits), citrus fruits (pulp), tobacco, sugar beet, red beet - 0.1; tea - 0.5*; wild berries and mushrooms - 0.01
456	Fencapton	Fruits (seeded fruits) - 0.3
457	Phenmedipham	Sugar beet, red beet - 0.2; chicory, endive - 0.5
458	Fenoxaprop-P- ethyl	Cereal grain, carrot, Red beet, sunflower (oil), onion - 0.01; sugar beet, soya (beans, oil) - 0.1; cabbage , sunflower (seeds)- 0.02; rape (grain, oil), pea - 0.2
459	Fenoxcarb	Grapes - 0.1; fruits (seeded fruits, stone fruits)-

		0.01
460	Derivatives of phenoxy-propanoic acid; Metabolites and half-products of synthesis of Centaur:	Sugar beet -0.02
	-2, 3, 5-trichloro-pyridine	NR
	-2-etoxy-ether 2-chloropropionic acid	NR
	-4-(3', 5'- dichloropyridil -2-oxy) phenol	NR
461	Fenpiclonil	NR
462	Fenpyroxymate	Soya (beans, oil), grapes, Fruits (seeded fruits, including apples)-0.3; Cattle kidney, liver - 0.02 ^{*,**} ; Cattle milk - 0.005 ^{*,**} ; hop (dry) - 10.0 ^{*,**} ; oranges (including hybrids) - 0.02 ^{*,**} ; Fruits (seeded fruits)-0.2; grapes-0.3
463	Fenpropatrine	Fruits (seeded fruits), grapes – 5.0; cottonseed (refined oil)-0.03 [*] ; bovine meat - 0.5 ^{*,**} ; bovine milk - 0.1 ^{*,**} ; bovine by-products - 0.05 ^{*,**} ; cottonseed (seed), tomatoes, sweet pepper (including bay pepper) - 1.0 ^{*,**} ; non-refined cottonseed oil - 3.0 ^{*,**} ; egg plants, pickling - 0.2 ^{*,**} ; eggs, poultry by-products - 0.01 ^{*,**} ; poultry meat- 0.02 ^{*,**} ; chili pepper (dry) - 10.0 ^{*,**} ; tea (green, black) - 2.0 ^{*,**} ; Fruits (seeded fruits), grapes-0.02; cottonseed (oil)-0.03*
464	Fenpropidin	Cereal grain - 0.25
465	Fenpropimorph	Cereal grain - 0.5 [*] ; sunflower (seeds) - 0.05 [*] ; sunflower (oil) - 0.1 [*] ; bananas - 2.0 ^{*,**} ; eggs, mammals fat (other than dairy fat), milk, fat, poultry meat and by products - 0.01 ^{*,**} ; bovine, goat, sheep, swine liver, sugar beet - 0.05 ^{*,**} ; bovine, goat, sheep, swine liver - 0.03 ^{*,**} ; mammals meat (other than sea mammals) - 0.02 ^{*,**} ; Cereal grain-0.2* ;
466	Fenthion	Cherries – 2.0 ^{*,**} ; citrus - 2.0 ^{*,**} ; olives, olive oil - 1.0 ^{*,**} ; shelled rice - 0.005 ^{*,**} ; cereal grain, grain legumes, sugar beet-0.15; milk and milk products-0.01; meat and meat products - 0.2
467	Fentoate	Citrus fruits (pulp) - 0.05 [*] ; berries-0.01; fruits (seeded fruits), grapes -0.1; cereal grain, rice, fruits (stone fruits) -0.1 [*]
468	Fenuron	Wild berries, mushrooms - 1.0
469	Fipronil	Potatoes – 0.02, cereal grain – 0.005; bananas - 0.005 ^{*,**} ; sunflower(seeds) - 0.02 ^{*,**} ; head cabbage, bovine kidneys and milk, eggs, poultry by-products, cabbage (including

		broccoli, Chinese and cauliflower), bovine liver - 0.1 ^{*,**} ; potatoes, cereal grain - 0.005, bovine meat - 0.5 ^{*,**} ; corn, poultry meat, rice - 0.01 ^{*,**} ; sugar beet - 0.2 ^{*,**} ;
470	Flamprop- izopropyl	Cereal grain - 0.1 *
471	Flamprop –M-methyl	Cereal grain- 0.06*
472	Florasulam	Cereal grain -0.05; corn (grain, oil) – 0.1
473	Fluazinam	Potatoes -0.025; seeded fruits, grapes – 0.05*
474	Fluazifop-P- butyl	Red beet -0.1; sugar beet, onion , potatoes - 0.02; carrot, pea - 0.03; fruits (seeded fruits, stone fruits) grapes- 0.02*; cabbage, rape (grain, oil) - 0.04; sunflower (oil, seeds), soya (beans, oil)-0.04
475	Fludioxonil	Cereal grain – 0.05; corn (grain) -0.02; sunflower (seeds, oil), sugar beet, potatoes, soya (beans, oil), rape (grain, oil)- 0.05; grapes (berries, juice))-2.0; peas (including green peas) – 0.3; apple meal dry - 20.0 ^{*,**} ; basil, green onion, head salad, mustard leaf, cress-salad - 10.0 ^{*,**} ; basil, green onion (dry) - 50.0 ^{*,**} ; black currant, blueberry (including boysenberry and loganberry), fruits seeded (other than pear), red and black raspberry - 5.0 ^{*,**} ; blueberry, head cabbage - 2.0 ^{*,**} ; broccoli, carrot, pear - 0.7 ^{*,**} ; citrus - 7.0 ^{*,**} ; cottonseed (seeds), eggs, mammals and poultry by-products - 0.05 ^{*,**} ; cucumbers, egg plants, pumpkin, legumes (other than feed and soya beans) - 0.3 ^{*,**} ; kiwi - 15.0 ^{*,**} ; poultry and mammals meat (other than sea mammals) milk, sweet corn (boiled in cobs) -0.01 ^{*,**} ; melon - 0.03 ^{*,**} ; bulb onion, tomatoes - 0.05 ^{*,**} ; sweet pepper (including bay pepper) - 1.0 ^{*,**} ; pistachio - 0.2 ^{*,**} ; strawberries - 3.0 ^{*,**} ; cereal grain, corn (grain) - 0.02; sunflower (seeds, oil), peas, sugar beet, potatoes, soya (beans, oil), rape (grain, oil) - 0.05; grapes (berries, juice) - 2.0
476	Flumetrine	Bovine meat - 0.2 ^{*,**} ; bovine milk - 0.05 ^{*,**} ;
477	Flumetsulam	Cereal grain -1.0
478	Flumioxazin	Sunflower (seeds, oil), soya (beans, oil) – 0.1
479	Fluometuron	Cottonseed (oil) - 0.1; Cereal grain -0.5*
480	Fluoxastrobine	Cereal grain – 0.5
481	Fluopicolide	Potatoes-0.05; Brussels cabbage - 0.2 ^{*,**} ; dry grapes (raisins), Wales onion - 10.0 ^{*,**} ; mammals by-products, mammals meat (other

		than sea mammals), poultry meat and by-products, eggs - 0.01 ^{*,**} ; cabbage (including broccoli, Chinese and cauliflower) - 2.0 ^{*,**} ; edible vegetable yields, pumpkins - 0.5 ^{*,**} ; grape meal, chili pepper (dry), grapes - 2.0 ^{*,**} ; milk - 0.02 ^{*,**} ;
482	Fluopiram	Cereal grain – 0.1; grape – 0.2 ^{**} ; fruits (seeded fruits) – 0.5 ^{**} ; fruits (stoned fruits) – 0.3 ^{**} ; bananas – 0.1. ^{**} ; tomatoes – 0.5 ^{**} ; pepper – 0.5 ^{**} , nuts – 0.3 ^{**} ; berries (strawberries and others) – 0.2 ^{**} ; cucumbers – 0.5 ^{**}
483	Fluroxypyr	Cereal grain, onion - 0.05
484	Flurochloridon	Cottonseed (oil)- 0.01; potatoes, sunflower (seeds, oil), carrot – 0.1;
485	Flusilasol	Apple and grape meal dry, mammals by-products - 2.0 ^{*,**} ; apricots, nectarines, peach, cereal grain, poultry meat and by-products - 0.2 ^{*,**} ; bananas - 0.03 ^{*,**} ; dry grapes (raisins), fruits seeded - 0.3 ^{*,**} ; eggs, rape (grain), soybean oil refined, sunflower (seeds) - 0.1 ^{*,**} ; mammals meat (other than sea mammals) - 1.0 ^{*,**} ; milk, soya (beans), sugar beet - 0.05 ^{*,**} ; sweet corn (boiled in cobs) - 0.01 ^{*,**} ;
486	Flutalonil	Eggs, mammals meat (other than sea mammals) milk, poultry meat and by-products - 0.05 ^{*,**} ; bovine, goat, swine, sheep kidneys - 0.1 ^{*,**} ; bovine, goat, swine, sheep liver - 0.2 ^{*,**} ; non-processed rice bran - 10.0 ^{*,**} ; rice shell out - 2.0 ^{*,**} ; milled rice - 1.0 ^{*,**} ;
487	Flutriafol	Cereal grain, corn (grain), millet, rice, pea, fruits (seeded fruits), sunflower (seeds, oil), grapes -0.05 sugar beet - 0.1; rape (grain, oil) – 0.2
488	Flufenzine	Fruits (seeded fruits)-0.04*, grapes-0.02*
489	Flucithrinat	Cereal grain -0.005
490	Fozalone	Cabbage, melons- 0.2*; cottonseed (oil), egg-plants, tomatoes, sugar beet, fruits (seeded fruits, stone fruits), grapes, citrus fruits (pulp), cereal grain, tobacco, mushrooms, grain legumes -0.2; potatoes, soya (beans, oil), oil poppy - 0.1; dry hop - 2.0*; rice - 0.3; livestock products, wild berries -0.01
491	Foxim	Cereal grain, Russian turnip, turnip, peas, sunflower (oil), corn (grain)- 0.05*; potatoes, tomatoes, egg-plants, meat - 0.02; cabbage,

		sugar beet - 0.1; sunflower (seeds) - 0.1 *; dry hop -0.5*; carrot, eggs – 0.01, Cereal grain after treatment under storage conditions - 0.6
492	Folpet	Potatoes – 0.1; grapes, fruits (seeded fruits) – 10.0, fruits (stone fruits)-0.02; cucumbers, bulb-onion - 1.0 ^{*,**} ; dry grapes (raisins) - 40.0 ^{*,**} ; salad (headed) - 50.0 ^{*,**} ; melon, tomatoes - 3.0 ^{*,**} ; strawberries - 5.0 ^{*,**} ;
493	Foramsulfuron	Corn (grain) -1.0; corn (oil)-0.5
494	Foreite	Dry beans, coffee beans, legumes (pods and/or unripe seed), cottonseed (seeds), corn, corn flour, soybeans, sorghum, sugar beet - 0.05 ^{*,**} ; corn oil non-refined - 0.1 ^{*,**} ; edible corn oil - 0.02 ^{*,**} ; potatoes - 0.2 ^{*,**} ; mammals meat and by-products (other than sea mammals) - 0.02 ^{*,**} ; meat, eggs - 0.05 ^{*,**} ; milk - 0.01 ^{*,**} ;
495	Formothion	Cottonseed (oil), sugar beet, red beet, fruits (seeded fruits, stone fruits), cabbage, grapes, tea, pomegranates - 0.2; citrus fruits (pulp) - 0.04*; dry hop - 2.0*
496	Fosmet	Sugar beet - 0.25 ; mushrooms - 0.1 ; wild berries – 0.01; potatoes – 0.05; blueberry, grapes, apricot, nectarine, peach, fruits seeded - 10.0 ^{*,**} ; citrus fruits - 3.0 ^{*,**} ; cottonseed (seed) - 0.05 ^{*,**} ; tree nuts - 0.2 ^{*,**} ; bovine meat - 1.0 ^{*,**} ; milk - 0.02 ^{*,**} ;
497	Ether phosphate (adjuvant)	RNR
498	Phosphine	Cereal grain - 0.1; grain products, sugar, dry vegetables and fruit, cacao beans, tea, spices, nuts, peanut-0.01; soya (beans)-0.05*
499	Fluorglycophen	Cereal grain – 0.01
500	Furathiocarb	Cereal grain, sunflower (seeds), rape (grain), corn (grain), sugar beet –0.02
501	Heptenophos	Cereal grain, grain legumes, fruits (seeded fruits, stone fruits), grapes, cucumbers, tomatoes, pepper - 0.1*; citrus fruits (pulp) - 0.05*; berries -0.01; potatoes - 0.01*
502	Quizalofop-P- ethyl	Red beet - 0.01; water melon, cabbage, onion, sugar beet, carrot, potatoes, tomatoes, rape (grain, oil) -0.05; soya (beans, oil), sunflower (seeds, oil) - 0.1; pea -0.4
503	Quinometionate	NR
504	Chloramben	Cabbage, tomatoes, grapes, citrus fruits (pulp), soya (beans, oil), cottonseed (oil) - 0.25
505	Chlorantraniliprol	Celery – 7.0 ^{*,**} ; cereal grain – 0.02 ^{*,**} ; cottonseed (seeds) – 0.3 ^{*,**} ; eggs – 0.01 ^{*,**} ;

		vegetables with edible fruits (except pumpkin, cucumbers, pepper, tomato) – 0.6*,**; pepper – 1.0*,**; cucumber – 0.3**,; tomato, eggplants – 0.6**,; pumpkin – 0.3**,; grapes – 1.0**,; raisin – 2.0**,; leaf vegetables (parsley and other) – 20.0*,**,salad (all types), cabbage (all types) – 20.0**,; citrus fruits – 1.0**,; meat of mammals (except sea), offal of mammals, milk, meat, poultry offal – 0.01*,**,; milk fat – 0.1*,**,; pepper Chili (dry) – 5.0*,**,; fruits (stone type) – 1.0**,; fruits seed type – 0.5; vegetables with edible roots and tubers – 0.02*,**,; potato – 0.1 Fruits (seeded fruits) – 0.5; potatoes – 0.1
506	Chlorbromuron	Cereal grain, corn (grain), soya (beans, oil) - 0.1; carrot – 0.2
507	Chlordane	Nuts (pecan, hazel nut, walnut) – 0.02*,**,; cottonseed oil, flax oil, soybean oil (crude) – 0.05*,**,; soybean oil, refined – 0.02*,**,; fruits and vegetables – 0.02*,**,; corn, rice (polished), sorghum, grain of cereals, eggs – 0.02*,**,; meat of mammals (except sea – control on fat) – 0.05*,**,; milk – 0.002*,**,; poultry meat (control on fat) – 0.5*,**
508	Chloridazon	Sugar beet, red beet-0.1
509	Chlormequat	Barley, wheat, rye – 2.0*,; cottonseed seeds – 0.5*,**,; eggs – 0.1*,**,; goat meat – 0.2*,**,; kidney of Cattle, goats, pigs, sheep – 0.1*,**,; meat of Cattle, pigs, sheep – 0.2*,**,; milk of Cattle, goat, sheep – 0.5*,**,; oats – 10.0*,**,; poultry meat – 0.04*,**,; poultry offal – 0.1*,**,; rapeseed (seeds) – 5.0*,**,; rapeseed oil, crude – 0.1*,**,; rye bran – 10.0*,**,; rye flour – 3.0*,**,; rye flour, not screened – 4.0*,**,; triticale – 3.0*,**,; wheat flour – 2.0*,**,; grapes, fruits (seed type), tomato, cabbage – 0.05
510	Chlorimuron-ethyl	Soya (beans, oil)- 0.05
511	Chlorinat	Cereal grain, vegetables (other than potato), fruits (seeded fruits, stone fruits) - 0.1
512	Chlormequat chloride	Cereal grain - 0.1; grapes, fruits (seeded fruits), tomatoes, cabbage - 0.05
513	Chlor-oxurone	Carrots – 0.02
514	Chlorothalonil	Tomato – 2.0; grapes – 0.5*,; cucumbers – 5.0*,; potato – 0.2; fruits (seed type) – 0.15. cereal grain – 0.1; hop (dry) – 1.0*,; beans (dry

		beans) – 0.2*,**; cabbage: broccoli, Brussels sprouts – 5.0*,**; cabbage (headed), cauliflower – 1.0*,**; carrot – 1.0*,**; celery – 10.0*,**; celery (leaf) – 3.0*,**; beans (pods or/and not ripened seeds) – 5.0*,**; onion (bulb) – 0.5*,**; parsley – 3.0*,**; fruits (stone type): peach – 0.2*,**; cherry – 0.5*,**; melon – 2.0*,**; banana – 0.01*,**; pumpkin – 5.0*,**; sweet corn (boiled cobs) – 0.01*,**; sugar beet – 0.2*,**; cranberry – 5.0*,**; pepper sweet, including clove pepper) – 7.0*,**; Chili pepper (dry) – 70.0*,**; peanut – 0.05*,** tomatoes – 0.15* ; fruits (seeded fruits), grapes – 0.15 ; cucumbers – 0.1* ; dry hop – 1.0* ; potatoes – 0.05 ; cereal grain – 0.1
515	Chlorpyrifos	Corn (grain), sugar beet, rapeseed (seed, oil) – 0.05; cottonseed oil for human consumption – 0.05*; cereal grain - 0.5; seed type fruits, grapes – 0.5; potato – 2.0; fruits (stone type) (except peach and nectarine) – 0.5**, peach, nectarine – 0.2**, citrus fruits – 0.3**, cabbage headed – 1.0**, almond, cauliflower, coffee beans, pecan, walnuts – 0.05*,**; bananas, broccoli, pepper sweet (including clove pepper), tea green and black - 2.0*,**; carrot, soya beans, wheat flour, dried grapes (raisin) – 0.1*,**; kidney, liver of Cattle., pig offal, beans (in pods and/or not ripened), eggs, green peas, poultry meat and offal, sheep offal, corn sweet (table, boiled in cobs – 0.01*,**, meat of Cattle and sheep, Chinese cabbage, cranberry – 1.0*,**; cottonseed (seed), strawberry – 0.3*,**; corn oil, onion (bulb)*,**; milk of Cattle., goat and sheep, pig meat - 0.02*,**; pepper Chili (dry) – 20.0*,**; rice, sorghum – 0.5*,**; soybean oil refined – 0.03*,** corn (grain) – 0.0006* ; rape (grain, oil) – 0.05 ; cottonseed (oil) – 0.0005* ; cereal grain – 0.01 ; fruits (seeded fruits) – 0.5 ; grapes – 0.4 ; potatoes, sugar beet – 0.005 ; fruits (stone fruits) – 0.2** ; citrus fruits – 0.3**
516	Chlorpyrifos-methyl	Meat, fat and offal of Cattle., and chicken – 0.005*,**; citrus fruits – 2.0*,**; eggplants, grapes, pepper, fruits seed type, tomato – 1.0*,**; Chili pepper (dry), sorghum, wheat (grain) – 10.0*,**; potato – 0.01*,**; rice –

		0.1*,**; stone type fruits – 0.5*,**; strawberry – 0.06*,**; wheat bran, not processed – 20.0*,**
517	Chlorpropham	Meat of Cattle – 0.1*,**; Cattle offal – 0.01*,**; milk fat – 0.02*,**; milk – 0.01*,**; potato – 30.0*,**; onion, carrot, chicory - 0.05; peeled potatoes for chips production-3.0
518	Chlorsulfoxym	Cereal grain, flax (oil), corn (corn) -0.005
	2-amine-4-dimethylamine-6-isopropylidene aminoxy-1,3,5-triazine, metabolite and half-product of synthesis of Krug	NR
519	Chlorsulfoxym - methyl	Cereal grain, corn (grain)- 0.005
520	Chlorsulfuron	Flax (seeds)-0.01; Cereal grain -0.01
	2-amine-4-methyl-6-metoxy-1,3,5-triazine, metabolite and half-product of synthesis of Hardin	NR
521	Potassium salt of chlorsulfuron	Flax (seeds) – 0.01
522	Chlortaldimethyl	potatoes- 0.002; vegetables, fruits (seeded fruits, stone fruits), fish, meat, butter – 0.05; milk products -0.04; sugar -0.02
523	Chlortholuron	Cereal grain - 0.0 1 *
524	Chlorphenetol	Cottonseed (oil), grapes -0.1*; citrus fruits (pulp) -0.1; fruits (seeded (fruits))-2.0
525	Chlorfluazuron	potatoes, cottonseed (oil) - 0.05
526	Cyanofos	Citrus fruits (pulp) - 0.05*; beet, cabbage, fruits (seeded fruits), grapes - 0.1
527	cyhalothrin	Almond, in shell – 2.0*,**; apricot, nectarine, peach, barley, cabbage (broccoli, Chinese, cauliflower) – 0.5*,**; asparagus, corn – 0.02*,**; berries and other small fruits, citrus, mango, vegetables with edible bulbs, kidney of Cattle, goats, pigs and sheep, milk, legumes, seeds of oilseeds, plums, fruits seed type – 0.2*,**; cabbage, dry grapes (raisin), vegetables with edible fruits (except pumpkin type), cherry – 0.3*,**; vegetables with edible fruits pumpkin type, liver of Cattle, goat, pig and sheep, oat, legumes, rye, sugar cane, triticale, wheat – 0.05*,**; meat of mammals (except sea), Chili pepper (dry) – 3.0*,**; olives, rice – 1.0*,**; vegetables with edible roots and tubers, tree nuts – 0.01*,**; wheat bran, not processed -0.1*,**
528	Cyhexatin	Cottonseed (oil), fruits (seeded fruits), grapes, citrus fruits (pulp) - 0.01; soya (beans, oil) -0.1

		; dry hop - 1.0
529	Cycloate	Sugar beet, red beet - 0.3
530	Cycloxydim	Beans (dry) – 2.0*,**; cabbage (head type, cauliflower) – 2.0*,**; carrot – 0.5*,**; beans ordinary (in pods and/or not ripened) -1.0*,**; grapes 0.5*,**; salad headed and leaf – 0.2*,**; peas (pods and seeds) – 1.0*,**; peas, shelled, juicy seeds – 2.0*,**; potato – 2.0*,**; rapeseed (seeds) – 2.0*,**; soybeans (beans, dry) – 2.0*,**; strawberry – 0.5*,**; sugar beet – 0.2*,**
531	Cymoxanil	Potatoes, cucumbers-0.05; grapes, tomatoes-0.1; sunflower (seeds, oil)-0.2; onion – 0.5
532	Zineb	Potatoes - 0.1; cereal grain, rice, pea -0.2; tomatoes, cucumbers, sugar beet, onion, gourds, fruits (seeded fruits, stone fruits), grapes- 0.6; dry hop, tobacco, essential oil rose -1.0; berries – 0.02
533	Cinidon-ethyl	NR
534	Aaphytora and ethylene thiuram disulfide (complex), metiram (synonym)	All food products - 0.02
535	Aaphytora and ethylene thiuram disulfide and manganese ethylene-bis-dithiocarbamate (blend)	Potatoes, fruits (seeded fruits), grapes - 0.1
536	Cypermethrin (ζ- and β- Cypermethrines)	Alfalfa - 30.0***; artichoke - 0.1***; barley, wheat, oat, rye – 2.0*; cabbage headed – 1.0*; carambola - 0.2***; grain cereals (other than barley, wheat, oat, rye) - 0.3***; citrus fruit – 2.0*; coffee beans - 0.05***; dray grapes (all kinds of raisin) - 0.05***; durian - 1.0***; egg plant - 0.03***; eggs – 0.01; fruit bearing vegetables other than pumpkin - 0.07***; grapes – 0.2, leaf vegetables - 0.7***; onion (leek, turnips) – 0.05; legumes - 0.7***; litchee - 2.0***; longan - 1.0***; mango - 0.7***; mammals meat (other than sea mammals) – 2.0*; dairy fat - 0.5***; milk – 0.05; oilseeds – 0.1***; okra - 0.05***; olive oil, refined - 0.5***; olive oil, non refined - 0.5***; olives - 0.05***; papaya - 0.5***; chili pepper – 2.0***; chili pepper dry - 10.0***; sweet pepper, including clove pepper – 0.1*; seeded fruits (including small fruits) – 0.7*; poultry meat - 0.1***; poultry by-products, edible - 0.05***; legumes - 0.05***; rice - 2.0***; root and bulb vegetables (other than sugar beets) - 0.01***;

		stone fruits – 2.0*; strawberries - 0.07 ^{*,**} ; sugar beet – 0.1*; sugar cane - 0.2 ^{*,**} ; sweet corn (boiled in cob) - 0.05 ^{*,**} ; tea (green, black fermented, dry) - 20.0 ^{*,**} ; tomatoes – 0.2; wheat bran unprocessed - 5.0 ^{*,**} ; cottonseed (oil) - 0.01*; sunflower (seeds, oil), gourds, cucumbers – 0.2; berries - 0.01; fish - 0.0015; pea, rape (oil), soya (oil), cultured mushrooms -0.1; potatoes, carrot, soya (beans), corn (grain) - 0.05; meat, livers and kidneys of cattle, sheep, pigs, poultry, fats - 0.2;
537	Cyprodinil	Fruits (seeded fruits, other than apples) – 1.0; apples – 0.05; stoned fruits – 2.0; grapes -5.0; carrot – 2.0 ^{*,**} ; almond in shell - 0.05 ^{*,**} ; almond - 0.02 ^{*,**} ; barley - 3.0 ^{*,**} ; legumes 9 other than feed and soya beans), sweet pepper(including clove pepper), raspberry, tomatoes, wheat - 0.5 ^{*,**} ; cucumbers, egg plants, pumpkin – 0.2 ^{*,**} ; dray grapes (raisins), prunes - 5.0 ^{*,**} ; mammals by-products, eggs, mammals meat (other than sea mammals), poultry meat and by-products - 0.01 ^{*,**} ; head salad and leaf salad - 10.0 ^{*,**} ; milk - 0.0004 ^{*,**} ; bulb-onion - 0.3 ^{*,**} ; strawberries, wheat bran unprocessed - 2.0 ^{*,**} ;
538	Cyproconazole	Cereal grain - 0.05; sugar beet, fruits (seeded), grapes-0.1
539	Cyprosulphamide	Corn (grain, oil) – 0.1
540	Cyromazine	Artichoke - 3.0 ^{*,**} ; dry beans - 3.0 ^{*,**} ; broccoli - 1.0 ^{*,**} ; celery - 4.0 ^{*,**} ; cucumbers - 2.0 ^{*,**} ; mammals by-products, edible - 0.3 ^{*,**} ; eggs - 0.3 ^{*,**} ; fruit bearing vegetables, other than pumpkin - 1.0 ^{*,**} ; salad, leaf and headed - 4.0 ^{*,**} ; lima bean (green pods and/or unripe beans) - 1.0 ^{*,**} ; mango - 0.5 ^{*,**} ; mammals meat (except for sea mammals) - 0.3 ^{*,**} ; melons, other than water melons - 0.5 ^{*,**} ; milk – 0.01 ^{*,**} ; mushrooms - 7.0 ^{*,**} ; leaf mustard - 10.0 ^{*,**} ; bulb-onion - 0.1 ^{*,**} ; chili pepper dry - 10.0 ^{*,**} ; poultry meat - 0.1 ^{*,**} ; poultry by-products - 0.2 ^{*,**} ; fruit-bearing vegetables other than pumpkin -3.0 ^{*,**} ; pumpkin - 2.0 ^{*,**} ;
541	Cyflutrine	Seeded fruits - 0.1 ^{*,**} ; cauliflower, citrus pulp (dry) - 2.0 ^{*,**} ; cottonseed (seeds) - 0.7 ^{*,**} ; cottonseed oil crude, mammals meat (other than sea mammals), chili pepper dry -1.0 ^{*,**} ;

		egg plants, pepper, tomatoes - 0.2 ^{*,**} ; potatoes, eggs, poultry meat and by-products - 0.01 ^{*,**} ; bovine, goat, swine, sheep kidneys, bovine, goat, swine, sheep liver - 0.05 ^{*,**} ; milk - 0.04 ^{*,**} ; rape (grain) - 0.07 ^{*,**} ;
542	Cyhexatine	Seeded fruits (apples, pears) - 0.2 ^{*,**} ; currant 9 red, black, white) - 0.1 ^{*,**} ; grapes - 0.3 ^{*,**} ; oranges 9 including hybrids) - 0.2 ^{*,**} ; chili pepper dry - 5.0 ^{*,**} ;
543	Edil	Potatoes, soya (beans, oil), sunflower (seeds, oil) -0.02
544	Emamectin benzoate	Grapes-0.05; cabbage-0.7; tomatoes-0.02
545	Endosulfan	Avocado, papaya, mango, pumpkin - 0.5 ^{*,**} ; tomatoes – 0.5; cocoa beans, coffee beans - 0.2 ^{*,**} ; cottonseed (seeds) - 0.3 ^{*,**} ; cucumbers – 1.0; egg plats - 0.1 ^{*,**} ; nuts (hazelnuts, macadamias - 0.02*; litchee - 2.0 ^{*,**} ; melon - 2.0 ^{*,**} ; potatoes, sweet potato - 0.05 ^{*,**} ; tea - 30.0 ^{*,**} ; eggs - 0.03 ^{*,**} ; mammals meat (other than sea mammals) - 0.2 ^{*,**} ; mammals kidneys - 0.03 ^{*,**} ; mammals liver - 0.1 ^{*,**} ; milk - 0.01 ^{*,**} ; dairy fat - 0.1 ^{*,**} ; poultry (meat and by-products) - 0.03 ^{*,**} ; soya (beans) - 1.0 ^{*,**} ; soya (oil) - 2.0 ^{*,**} ; apple crème - 0.5 ^{*,**} ; berries – 0.002; eucumbers, tomatoes-0.002; cottonseed (oil)-0.05 cottonseed (oil)-0.05
546	Endrine	Vegetables with edible yields, pumpkins - 0.05 ^{*,**} ; poultry meat - 0.1 ^{*,**} ;
547	Epoxyconazole	Cereal grain-0.2; sugar beet –0.05
548	Esfenvalerate	Eggs – 0.01 ^{*,**} ; poultry meat and offal – 0.01 ^{*,**} ; corn (grain) -0.01*; sunflower (seeds), soya (beans) -0.02*; sunflower (oil), soya (oil) –0.04*; sugar beet – 0.01*; cottonseed (oil), potatoes, grapes, peas, cereal grain, fruits (seeded fruits), rape- 0.1; cabbage - 0.05; meat and meat products, milk-0.01
549	Ethaboxam	Potatoes-0.5; grapes-3.0
550	Etalfluralin	Water melons - 0.05*; cottonseed (oil), sunflower (seeds, oil), soya (beans, oil) – 0.02
551	Ethefon	Fruits (seed type) – 5.0 ^{*,**} ; fruits (stone type) – 10.0 ^{*,**} ; cereal grain – 1.0*; blueberry – 20.0 ^{*,**} ; cantaloupe – 1.0 ^{*,**} ; eggs – 0.2 ^{*,**} ; cottonseed (seed) – 2.0 ^{*,**} ; raisin – 5.0 ^{*,**} ; figs – (dry, candied) – 10.0 ^{*,**} ; grapes – 1.0 ^{*,**} ; nuts: hazel – 0.2 ^{*,**} , walnuts – 0.5 ^{*,**} ; pepper – 5.0 ^{*,**} ; chili Pepper (dry) –

		50.0*,**; pineapple – 2.0*,**; meat (Cattle, Goat, horse, pigs, sheep) – 0.1*,**; offal (Cattle, goat, horse, pig, sheep) – 0.02*,**; milk (Cattle, sheep, goat) – 0.05*,**; poultry meat – 0.1*,**; poultry offal – 0.2*,**; tomato – 2.0*; citrus fruits, sugar beet, pea, cabbage, cucumbers – 0.5*; potato – 0.15 tomatoes 0.5*;
552	Ethylene thiourea	All plant and food products -0.02
553	Ethyl mercuric chloride (Granozane)	All food products and raw material – 0.005
554	Ethylfenacin	RNR
555	Ethyofencarb	potatoes - 0.04; grain legumes -0.2*; sugar beet - 0.1*; cottonseed (oil) , cereal grain, rice - 0.05*; dry hop - 1.0*
556	Ethirimol	Cereal grain - 0.05
557	Ethoxyquin	Peach – 3.0*,**
558	Aliphatic alcohol ethoxylate C ₈ -C ₁₀	NR
559	Isodecyl alcohol ethoxylate (adjuvant)	RNR
560	Ethoprophos	Strawberry, banana, sugar cane, melon – 0.02*,**; pepper, potato, sweet potato – 0.05*,**; tomato, cucumbers – 0.01*,**; Chili pepper (dry) – 0.2*,**; meat of mammals (except sea) – 0.01*,**; milk, offal of mammals – 0.01*,**; garden turnip – 0.02*,**
561	Etofenprox	Cottonseed (oil), potatoes - 0.1*; fruit (seeded fruits) – 1.0*,** fruits (seeded fruits) – 0.3*
562	Ethofumezate	Red beet, sugar beet -0.1; tobacco -1.0*
563	Etrimfos	Cottonseed (oil), fruits (seeded fruits, stone fruits), grapes -0.5*; sugar beet - 0.01*; cabbage, potatoes, sunflower (seeds, oil) -0.1*; pea, cereal grain (stored supplies) - 0.2*; berries (any) -0.01